




Emergency Medical Services Outcomes Evaluation



This publication is distributed by the U.S. Department of Transportation, National Highway Traffic Safety Administration, in the interest of information exchange. The opinions, findings and conclusions expressed in this publication are those of the author(s) and not necessarily those of the Department of Transportation or the National Highway Traffic Safety Administration. The United States Government assumes no liability for its contents or use thereof. If trade or manufacturers' names are mentioned, it is only because they are considered essential to the object of the publication and should not be construed as an endorsement. The United States Government does not endorse products or manufacturers.

Technical Report Documentation Page

1. Report No. DOT HS 809 603	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Emergency Medical Services Outcomes Evaluation		5. Report Date July 2003	
		6. Performing Organization Code	
7. Author(s) Ronald Maio, DO, MS		8. Performing Organization Report No.	
9. Performing Organization Name and Address University of Michigan Department of Emergency Medicine Ann Arbor, MI 48109-1382		10. Work Unit No. (TRAIS)	
		11. Contract or Grant No.	
12. Sponsoring Agency Name and Address National Highway Traffic Safety Administration Office of Research and Traffic Records Research and Evaluation Division 400 7 th Street, S.W., Washington, D.C. 20590		13. Type of Report and Period Covered	
		14. Sponsoring Agency Code	
15. Supplementary Notes			
16. Abstract The provision of prehospital (EMS) care has come under increased scrutiny in recent years. Many have questioned the value of the range of EMS services currently provided. There is a persistent concern about the lack of proof of effectiveness related to most EMS care. Clinical effectiveness studies to address EMS outcomes research require the development of sophisticated case-severity and effectiveness measures. Outcomes research will allow future generations of Americans to have an EMS system that provides both quality and cost-effective EMS care. This report describes a project that has laid the foundation for these clinical effectiveness studies to take place. This project developed a "blueprint" or "set of tools" that EMS practitioners can use to evaluate the effectiveness of EMS, or prehospital, care.			
17. Key Words EMS, Outcomes Research, clinical studies		18. Distribution Statement	
19. Security Classif. (of this report)	20. Security Classif. (of this page)	21. No. of Pages	22. Price

Table of Contents

Executive Summary.....	3
1.0 Introduction	6
2.0 Background.....	6
3.0 Conceptual Models for Prehospital Outcomes Research	7
4.0 Goals and Objectives of the Work Performed.....	9
5.0 Project Findings.....	10
5.1 Finding 1	10
5.2 Findings 2 and 3	19
5.2.1 Core Risk Adjustment Measures	19
5.2.2 Core Outcome Measures	23
5.2.3 Pain	27
5.2.4 Condition-Specific Risk Adjustment and Outcome Measures	28
5.2.4.1 Major/Minor Trauma	29
5.2.4.2 Respiratory Distress	36
5.2.4.3 Cardiac Arrest	41
5.2.4.4 Chest Pain	42
5.2.4.5 Respiratory Arrest/Airway Obstruction	42
5.2.4.6 Pediatric Seizures	42
5.3 Finding 4.....	42
5.4 Finding 5.....	43
6.0 Limitations.....	44
7.0 Implications	45
8.0 Future Research	46
9.0 Project Participants	47
10.0 List of Publications.....	48
11.0 List of Presentations	48
References	50

Appendices

Appendix I: ..Questionnaire respondents and their affiliations	63
Appendix II: ..Sample item from questionnaire.....	67
Appendix III: Measure identification process	69
Appendix IV: ..List of stakeholders.....	73

Executive Summary

Approach. The provision of prehospital care has come under increased scrutiny in recent years. Although it is acknowledged that timely transport is necessary for some patients, many have questioned the value of the range of prehospital care services currently provided. In the broader health care community, there is a persistent concern about the lack of proof of effectiveness related to most prehospital care. Most experts on both sides of the argument agree that methodologically sound outcomes research that identifies “what works” in prehospital care is long overdue. Clinical effectiveness studies to address EMS outcomes research require the development of sophisticated case-severity and effectiveness measures. Outcomes research will allow future generations of Americans to have an EMS system that provides both quality and cost-effective prehospital care.

Objectives. The primary goal of this work is to support and facilitate EMS outcomes research and evaluation to be conducted by the broad EMS community. In essence, a “blueprint” and a set of “tools” were developed that EMS practitioners can use to evaluate the effectiveness of prehospital care. The following objectives were met: 1) Identify conditions that should take precedence in EMS outcomes research; 2) Determine the appropriate risk adjustment measures for the priority conditions identified; 3) Determine the appropriate outcome measures for the priority conditions identified; 4) Identify important stakeholders and constituencies; 5) Develop a research dissemination plan.

Results. Seven conditions were identified for children and adults that should take precedence in EMS outcomes research. For adults, these conditions are minor trauma, respiratory distress, chest pain, major trauma, cardiac arrest, airway obstruction, and respiratory arrest. For children, these conditions are minor trauma, major trauma, respiratory

distress, airway obstruction, respiratory arrest, cardiac arrest and seizures. Next, core risk adjustment measures were identified that should be considered for all conditions. These include patient age, sex, race and ethnicity, vital signs, level of responsiveness, Glasgow Coma Scale (GCS), time intervals, and the EMS provider impression. Pain, using an adjective response scale (ARS) and a numeric response scale (NRS), should be measured for all patients. Condition-specific risk adjustment measures were identified for certain conditions. Outcome measures that should be considered for all conditions include survival, cost-effectiveness, and satisfaction. However, no measures that could be used as core measures for satisfaction or cost were identified. Outcome measures pertaining to disease progression, discomfort and dysfunction may also be condition-specific. Risk adjustment measures may also function as outcome measures depending upon the point in time during the episode of care the measure is applied. The following table provides an example using three conditions.

<u>Adult Priority Condition</u>	<u>Risk adjusters</u>	<u>Outcome measures</u>
Minor trauma	Revised Trauma Score, Glasgow Coma Scale (GCS); Abbreviated Injury Scale (AIS), Injury Severity Score (ISS), New ISS (NISS)	See risk adjusters SF-36
Respiratory distress	Peak Expiratory Flow Rate (PEFR), Pulse Oximetry; Visual Analog Scale for Dyspnea (discomfort)	See risk adjusters Mortality
Major trauma	Revised Trauma Score, GCS, AIS, ISS, NISS	See risk adjusters, SF-36, FCI, Mortality, Probability of Survival (Ps), Preventable Death Rate (PDR)

Conclusion. Priority conditions have been identified on which EMS outcomes research should be focused. Risk adjusters and outcome measures have been identified that

investigators can use as tools in EMS outcomes research. These should be evaluated in the prehospital setting. A particular focus of future research should be the identification, development and evaluation of satisfaction and cost outcome measures. Past and ongoing dissemination have included presentations and publications in peer-reviewed journals. Through all of these efforts, the ultimate goal is to conduct EMS outcomes research that will improve the delivery of prehospital care, and will thus benefit the general public.

1.0 Introduction

This paper is the final report for the Emergency Medical Services Outcomes Project (DTNH22-96-H-05245), for the period July 1, 1996 to March 31, 2002.

2.0 Background

The provision of prehospital care has come under increased scrutiny in recent years. Although it is acknowledged that timely transport is necessary for some patients, many have questioned the value of the range of prehospital care services currently provided (Callahan, 1997; Koenig, 1995, 1996; Reines et al., 1988; Smith et al., 1985; Spaite et al., 1995). In the broader healthcare community, there is a persistent concern about the lack of proof of effectiveness related to most prehospital care (Raskin, 1991; Relman, 1988; Roper et al., 1988). Most experts on both sides of the argument agree that methodologically sound outcomes research that identifies “what works” in prehospital care is long overdue (Callahan, 1997; Delbridge et al., 1998; NHTSA, 1996, Spaite et al., 1997; Brice et al., 1996; Spaite, 1993).


In 1994, NHTSA convened a workshop on methodologies for “measuring morbidity outcomes in EMS.” The experts in this workshop concluded that implementation of EMS outcomes research was essential (NHTSA, 1994). However, it was noted that the methods applicable to prehospital outcomes, especially those using non-mortality measures, had never been developed. These methods should be applicable across the entire spectrum of the “Six Ds” of patient outcomes: Death (survival); Disease (impaired physiology); Disability; Discomfort; Dissatisfaction and, Destitution (cost) [NHTSA, 1994]. In response to these conclusions, a five-year cooperative agreement, the Emergency Medical Services Outcomes

Project (EMSOP) was funded to facilitate EMS outcomes research and to implement the recommendations from the workshop of 1994.

3.0 Conceptual Models for Prehospital Outcomes Research

Development of methodologically acceptable outcomes models for EMS is long overdue. The EMSOP steering committee and consultants propose a conceptual framework that will provide a foundation for future EMS outcomes research using two distinct, conceptual models: 1) The “Episode of Care Model” (see Figures 1 and 2) [Spaite et al., 2001]; and 2) The “Prehospital Unit of Service Model” (see Figure 3) [Spaite et al., 2001].

**FIGURE 1
THE EPISODE OF CARE**



*Such as surgery, interventional radiology, etc.

**Such as specialty follow-up care, physical therapy, occupational therapy, etc.

FIGURE 2
THE EPISODE OF CARE MODEL
 Model for identifying the impact of treatment from each "unit of service"
 in the episode of care



PREHOSPITAL CARE	EMERGENCY DEPARTMENT CARE	EMERGENT SUBSPECIALTY CARE	INPATIENT CARE	FOLLOW-UP CARE
RA T OUT	RA T OUT	RA T OUT	RA T OUT	RA T OUT



RA = Risk Adjustment Measures
 T = Therapeutic intervention(s)
 OUT = Outcome Measures

FIGURE 3
PREHOSPITAL UNIT OF SERVICE MODEL
FOR EMS OUTCOMES RESEARCH

PREHOSPITAL CARE	EMERGENCY DEPARTMENT CARE	EMERGENT SUBSPECIALTY CARE	INPATIENT CARE	FOLLOW-UP CARE
------------------	---------------------------	----------------------------	----------------	----------------



*Modified from Spaite, et al: *Ann Emerg Med* 1993;22:639

The Episode of Care Model is useful in conditions where interventions and outcomes, especially survival and major physiologic dysfunction, are linked in an extremely time-dependent manner. Non-traumatic cardiac arrest is the prototypical condition for utilizing this model. The Prehospital Unit of Service Model is essentially a sub-unit of the Episode of Care Model. It is valuable for evaluating conditions that have minimal to moderate therapeutic time-dependency. This model should be used when one is most concerned about studying outcomes limited to the prehospital interval. An example of an outcome that could be studied using this model is pain from injuries sustained in a motor vehicle crash. These models should be broadly applied to a wide spectrum of conditions and interventions. We believe that these will be particularly useful in the evaluation of major trauma patients. Further, these models can be applied across the entire spectrum of the “Six Ds” of patient outcomes.

4.0 Goals and Objectives of the Work Performed

The primary goal of this work is to support and facilitate EMS outcomes research and evaluation to be conducted by the broad EMS community. In essence, a “blueprint” and a set of “tools” has been developed that EMS practitioners can use to evaluate the effectiveness of prehospital care. Objectives critical to the development of an EMS outcomes research plan included:

1. Identify conditions that should take precedence in EMS outcomes research;
2. Determine the appropriate risk adjustment measures for the priority conditions identified;
3. Determine the appropriate outcome measures for the priority conditions identified;
4. Identify stakeholders and constituencies important for EMS outcomes research;

5. Develop a research dissemination plan.

An objective identified at the start of the project, "identify existing data sets and evaluation methods useful for EMS outcomes research," was removed because the steering committee determined that there were no data sets in existence that could be used to actually conduct EMS outcomes research.

5.0 Project Findings

5.1 Finding 1: Identify conditions that should take precedence in EMS outcomes research

During phase one of this portion of the project, a list of EMS conditions was developed. An EMS condition was defined as an illness, injury or combination of signs and symptoms that caused EMS activation. A preliminary list of such conditions was identified using the NHTSA Uniform Data Conference data element items "Provider Impression" (data element 50), "Signs and Symptoms Present" (data element 52) and "Injury Site and Type" (data element 53) [NHTSA, 1993].

In phase two, frequency data were obtained for all the conditions identified. No local, state or federal databases were suitable for use due to inconsistent data definitions, inconsistent data formatting, and variation in inclusion criteria. EMS Data Systems, Inc. (Phoenix, Arizona) was selected to provide frequency data. EMS Data Systems collects data from various EMS systems across the country using optically-scanned data entry forms and data sets similar to that promulgated by NHTSA. Data from July 1, 1995 through June 30, 1996 were obtained from Alabama, Mississippi, Oklahoma, Illinois and eleven central California counties and used for the frequency analysis.

In phase three, the relevance of various outcomes and the potential impact of EMS on these outcomes, for each condition, was determined. Lacking meaningful outcome data, the investigators obtained expert opinions from 37 EMS researchers and leaders regarding the relevance and potential impact of EMS (see Appendix I). Respondents were asked to complete two questionnaires, one for patients less than 15 years of age and another for patients 15 years of age or older (see Appendix II). Two questions were included in the adult and child questionnaire: 1) For each of the following conditions, how would you rate the relevance of the following 6 outcome categories? and 2) For each of these conditions, how would you rate the potential impact of EMS (including both basic and advanced EMT care) on each outcome? The six outcome categories were defined as: survival (death), impaired physiology (disease), limit disability (disability), alleviate discomfort (discomfort), satisfaction (dissatisfaction), and cost (destitution). Respondents were asked to note the relevance and potential impact of EMS for each condition/outcome category on a 5-point scale ranging from (1) low to (5) high impact.

In phase four, a summary index score was developed and the conditions ranked according to this index. For each condition, a standard normal deviate (position of score on a standard normal distribution with mean of 0 and standard deviation of 1) was calculated. This transformation gives all variables the same mean and standard deviation. The frequency count and weighted score for each condition and age category were then multiplied together (in order to have positive signs after multiplication, negative z-scores were eliminated by the linear transformation of adding 4 to each score). Using this summary index score, the conditions were rank ordered for each age category.

The internal consistency of the total score (summed over the 6 Ds) for each condition was determined by calculating Cronbach's alpha. Correlation between impact and relevance scores was determined using the Spearman rank correlation coefficient. Agreement of rankings between respondents was measured in two different manners. First, respondents were divided into three, mutually exclusive, categories: 1) Physicians and non-physicians that were investigators or co-investigators; 2) Other physician respondents; and 3) Non-physician responders. Rankings were calculated within groups, then correlations among the three were calculated (Spearman rank correlation). Second, the average correlation among rankings by all respondents was calculated.

A sensitivity analysis was conducted by calculating index scores and ranking conditions using only survival scores and then using only discomfort scores. The rankings from each calculation were then compared to the original index score ranks that used all outcome categories, and Spearman rank correlation coefficients were calculated. See Tables 1 and 2.

Table 1. Ranking for priority conditions, children

<u>Condition</u>	<u>%Freq</u>	<u>Weighted Score</u>	<u>SD*</u>	<u>Index Score</u>
Minor trauma	51.3	9.6	4.52	34.3
Major trauma	7.9	17.1	4.79	25.7
Respiratory distress	10.0	15.0	5.00)	24.4
Airway obstruction	1.1	17.9	4.58	22.6
Respiratory arrest	.4	16.3	5.02	20.6
Cardiac arrest	.8	14.4	4.88	19.1
Seizure	14.1	9.0	3.49	19.0
Shock	< .1	13.9	4.41	18.2
Allergic reaction	.5	13.1	4.77	17.7
Environmental exposure	.6	12.2	4.49	17.0
Diabetes complication	.3	12.2	5.29	16.8
Cardiac problem	.2	11.2	5.14	15.9
Poisoning/OD	3.0	8.9	3.57	14.7
Hemorrhage	< .1	10.0	4.28	14.6
Chest pain	.9	9.4	5.47	14.4
Altered LOC	1.5	8.7	3.52	14.0
Fever	2.3	7.6	4.22	13.2

Table 1. Ranking for priority conditions, children (continued)

<u>Condition</u>	<u>%Freq</u>	<u>Weighted Score</u>	<u>SD*</u>	<u>Index Score</u>
Preg/labor/childbirth	.6	8.1	4.69	13.2
Stroke/CVA	< .1	7.6	5.05	12.5
Abdominal pain	1.2	6.7	3.50	12.0
Abdominal distress	1.4	5.9	3.22	11.3
Hypertension	<.1	6.1	3.56	11.1
Drug/alcohol problem	.2	5.8	3.24	10.9
Gyn problem	.1	5.7	3.68	10.7
Syncope/near syncope	.7	5.3	3.49	10.5
Dizziness	.2	4.5	2.84	9.7
Behavioral problem	.6	4.3	2.58	9.6

*SD=standard deviation

Table 2. Ranking for priority conditions, adults

<u>Condition</u>	<u>%Freq</u>	<u>Weighted Score</u>	<u>SD*</u>	<u>Index Score</u>
Minor trauma	36.1	10.3	4.64	33.3
Respiratory distress	13.0	15.3	4.82	27.3
Chest pain	10.2	14.8	4.56	24.8
Major trauma	3.6	17.1	4.86	22.3
Airway obstruction	.2	17.6	5.14	20.1
Cardiac arrest	2.2	15.9	5.57	20.1
Respiratory arrest	.2	16.3	5.22	18.9
Cardiac problem	3.3	13.3	4.39	18.3
Shock	.4	14.7	4.60	17.7
Diabetes complication	2.3	12.8	4.97	17.1
Allergic reaction	.4	13.8	4.90	16.9
Environmental exposure	.3	12.3	4.62	15.5
Stroke/CVA	2.7	10.3	5.31	15.0
Seizure	4.8	9.0	3.64	14.9
Altered LOC	3.7	9.1	4.17	14.4
Hemorrhage	.3	10.4	3.76	13.8
Poisoning/OD	1.8	9.0	3.80	13.4

Table 2. Ranking for priority conditions, adults (continued)

<u>Condition</u>	<u>% Freq</u>	<u>Weighted Score</u>	<u>SD*</u>	<u>Index Score</u>
Preg/labor/childbirth	1.0	9.4	3.97	13.3
Abdominal pain	4.1	7.2	3.72	12.7
Hypertension	1.0	7.9	3.35	11.9
Syncope/near syncope	1.8	7.0	4.64	11.4
Abdominal distress	2.9	6.4	3.23	11.4
Gyn problem	.3	6.7	3.06	10.6
Fever	.5	6.3	3.38	10.3
Drug/alcohol problem	.3	6.0	3.52	10.0
Dizziness	1.1	5.5	3.28	9.8
Behavioral problem	1.6	4.6	2.68	9.2

*SD=standard deviation

Following data analysis, EMSOP investigators and consultants met to determine what conditions were to be recommended as priorities for EMS. Criteria for selection were based on summary index scores, the proportion of EMS transports represented, and the feasibility of identifying risk adjustment measures and outcome measures for these conditions within the time and resource constraints of the project. For adults, these conditions (in order of importance) are minor trauma, respiratory distress, chest pain, major trauma, cardiac arrest, airway obstruction, and respiratory arrest. For children, these conditions (in order of importance) are minor trauma, major trauma, respiratory distress, airway obstruction, respiratory arrest, cardiac arrest and seizure. For adults, the top quartile conditions account for 65% of adult emergency transports and for children, 85% of emergency transports. Relief of discomfort is the outcome parameter EMS professionals identified as having the most potential impact for the majority of both adults and children in the top quartile conditions. EMS research priorities focusing on these conditions will ensure that scarce resources will be directed to conditions that not only affect a substantial portion of EMS patients, but also have the potential of providing the greatest benefit (see Tables 3 and 4) [Maio, 1999].

Table 3. Weighted score for top quartile conditions by outcome category, children

<u>Condition</u>	<u>Survival</u>	<u>Impaired Physiology</u>	<u>Limit Disability</u>	<u>Alleviate Discomfort</u>	<u>Satisfaction</u>	<u>Cost Effectiveness</u>
Minor Trauma	3.7	6.8	10.7	16.3	15.3	9.5
Major Trauma	20.1	18.1	19.0	16.3	16.7	14.2
Respiratory Distress	14.7	18.0	12.6	18.8	16.7	11.7
Airway Obstruction	24.3	20.1	18.4	16.9	17.3	13.4
Respiratory Arrest	23.5	21.0	20.4	10.5	13.8	13.0
Cardiac Arrest	21.2	20.0	19.0	5.9	13.6	12.5
Seizure	7.0	10.9	9.7	9.7	11.1	7.4

Table 4. Weighted score for top quartile conditions by outcome category, adults

<u>Condition</u>	<u>Survival</u>	<u>Impaired Physiology</u>	<u>Limit Disability</u>	<u>Alleviate Discomfort</u>	<u>Satisfaction</u>	<u>Cost Effectiveness</u>
Minor Trauma	4.1	6.8	12.2	17.2	15.6	11.0
Respiratory Distress	14.9	18.1	13.1	19.4	17.3	12.3
Chest Pain	14.2	14.3	12.1	20.5	17.6	12.5
Major Trauma	20.1	17.3	19.6	15.9	16.8	14.9
Cardiac Arrest	22.4	20.9	19.5	7.3	14.8	14.9
Airway Obstruction	24.0	19.3	18.6	18.1	16.4	12.4
Respiratory Arrest	23.7	20.4	20.7	11.0	13.4	13.6

5.2 Findings 2 and 3: Determine the appropriate risk adjustment measures for the priority conditions identified; Determine the appropriate outcome measures for the priority conditions identified.

5.2.1 Core Risk Adjustment Measures

Risk adjustment allows better judgment about the effectiveness and quality of alternative therapies; it fosters a comparison of apples with apples. By measuring risk adjustment measures (RAMs), researchers account for an important source of variation in their studies (Seidel et al., 1999). Core RAMs are necessary for prehospital outcomes research involving any EMS condition. Core RAMs should also be used, if possible, in retrospective EMS studies and for prospective prehospital investigations where applicable. The selection of the core RAMs by the EMSOP project team was based on four criteria: 1) they are “medically meaningful” for any EMS condition (ie, there is a possible connection between the RAM and any outcome pertinent to EMS); 2) data for the RAMs is readily available or measurable in the prehospital environment or may be obtained through linkage to other healthcare databases; 3) the RAMs have been described as risk attributes for outcomes research in other health settings; and 4) they were agreed to by all the EMSOP investigators and consultants.

The EMSOP investigators and consultants determined that the following core RAMs should be measured at the time of prehospital care: Patient age, sex, race and ethnicity, vital signs, level of responsiveness, GCS, time intervals, and the EMS provider impression (Zuckerman et al., 1998; Garrison et al., 2002; Donabedian, 1987; Fletcher et al., 1988; Shann et al., 1997; Pearson et al., 2001) [See Table 5].

Table 5. Recommended patient factors that are measured or collected at the time of out-of-hospital care that should be evaluated for use as core risk adjustment measures (RAMs) in out-of-hospital research.

RISK ADJUSTMENT MEASURE	VALUES
Age	Ideal: date of birth (DOB). If DOB unavailable, record in years unless an infant or toddler whose age should be recorded in months
Sex	Male or Female
Race and Ethnicity (Self-Reported)	White, non-Hispanic; White, Hispanic; Black, non-Hispanic; Black, Hispanic; American Indian or Alaskan Native; Or Asian or Pacific Islander
Initial Vital Signs	Pulse; Respiratory Rate; and Systolic Blood Pressure (BP),
Vital Signs Before and After a Major Intervention	Pulse; Respiratory Rate; and Systolic BP
Final Vital Signs (at time of transfer of care)	Pulse; Respiratory Rate; and Systolic BP
Initial Level of Responsiveness	AVPU: Alert, Verbal Response, Painful Response or Unresponsive
Level of Responsiveness Before and After an Intervention	AVPU
Final Level of Responsiveness (at time of transfer of care)	AVPU
Initial Glasgow Coma Scale (GCS) (Eye Opening + Motor Response + Verbal Response)	EYE OPENING None (1) To Pain (2), To Verbal Command (3) Spontaneous (4) MOTOR RESPONSE None (1), Abnormal Extension (Decerebrate) (2), Abnormal Flexion (Decorticate) (3), Withdrawal (Normal Flexion) (4), Localizes Pain (5), Obeys Commands (6) VERBAL RESPONSE None (1) Incomprehensible Sounds (2) Inappropriate Words (3) Confused Conversation (4) Oriented (5)

GCS Before and After an Intervention	As Above
Final GCS (at time of transfer of care)	As Above
Event to Treatment Interval	Number of Minutes from the Time when Incident Reported to the Time of Arrival of EMS at the Patient.
Prehospital Treatment Interval	Number of Minutes from the Time of Arrival of EMS at the Patient to the Time of Arrival at the Transport Destination
EMS Provider Impression of Presenting Condition	Abdominal Pain/Problems; Airway Obstruction; Allergic Reaction; Altered Level of Consciousness; Behavioral/Psychiatric Disorder; Cardiac Arrest; Cardiac Rhythm Disturbance; Chest Pain/Discomfort; Diabetic Symptoms (Hypoglycemia); Electrocution; Hyperthermia; Hypothermia; Hypovolemia/Shock; Inhalation Injury (Toxic Gas); Obvious Death; Poisoning/Drug Ingestion; Pregnancy/OB Delivery; Respiratory Arrest; Respiratory Distress; Seizure; Sexual Assault/Rape; Smoke Inhalation; Stings/Venomous Bites; Stroke/CVA; Syncope/Fainting; Traumatic Injury (+Cause of Injury Code); Vaginal Hemorrhage; Other; Unknown

The routine vital signs for the EMS patient are heart rate, respiratory rate, and blood pressure. Vital signs should be obtained and recorded as part of the initial assessment of every EMS patient. Since vital signs may also be used as an outcome measure, they should also be measured before and after an intervention and at the time of final EMS assessment. While there is no universally accepted method for assessing the level of responsiveness, the AVPU method (alert, verbal response, painful response, unresponsive) should be used. Since the

GCS may also be used as an outcome measure, it should also be measured before and after an intervention and at the time of final EMS assessment. The time intervals that should be measured as RAMs are the event-to-treatment interval (ETI) and the prehospital treatment interval (PTI). To calculate the ETI, you need the single point in time marking the start of the event (using the time the call is received at a public safety answering point) and the single point in time of the arrival of the EMS providers at the side of the patient. The PTI is calculated from the single point in time of arrival at the patient until single point in time that marks the arrival at the hospital.

The EMSOP investigators and consultants determined that other core RAMs should be obtained through linkage to other data sources, to include principal diagnosis and patient co-morbidity (see Table 6).

Table 6. Recommended core RAMs that should be obtained through linkage to other data sets.

RISK ADJUSTMENT MEASURE	VALUES
Principal Diagnosis	Primary ICD-9-CD Code on Uniform Billing Record
Patient Co-Morbidity	Secondary ICD-9-CD Codes on Uniform Billing Record

Principal diagnosis is a critical element in risk adjustment since risks may differ significantly depending on the diagnosis (Iezzoni, 1994). For most prehospital patients, the diagnosis used in risk adjustment for prehospital outcomes research will be the principal diagnosis received in the hospital. Patient co-morbidity includes all those conditions and complications that may put a patient with them at risk for a different outcome. A patient's co-morbidity can affect both short and long-term outcomes (Iezzoni, 1994).

5.2.2 Core Outcome Measures

In identifying core outcome measures, we used four criteria that were very similar to those used for identifying risk adjustment measures: 1) they are “medically meaningful” for any EMS condition; 2) data for the outcome measures are readily available or measurable in the prehospital environment or may be obtained through linkage to other healthcare databases; 3) the outcome measures have been used in outcomes research in other health settings; and 4) they were agreed to by all the EMSOP investigators and consultants.

The outcome categories identified for which specific outcome measures would be identified are the six “Ds”: death, disease (disease progression), disability, discomfort, destitution (cost), and dissatisfaction. Of note is that risk adjustment parameters that measure disease progression and discomfort can also be used as outcome measures and, therefore, core outcome measures. All of these core outcome measures can be measured or collected by EMS professionals in the prehospital setting.

The outcome parameter of mortality has been extensively used in EMS research and should continue to be used. The interval of care during which the patient was pronounced dead should also be obtained. At a minimum, mortality should be described using the following categories: death in the prehospital interval, death in the emergency department care interval, and death after hospitalization but before discharge. Death in intervals other than the prehospital interval will require data linkage outside of the direct control of prehospital systems. An outcome measure that is a crude indirect measure of disease severity and is related to mortality is patient disposition. Categories for this measure are treated/released at scene; treated and released from the emergency department; treated in the emergency department and admitted to hospital; discharged alive from hospital. As with

mortality, obtaining information for disposition will require data linkage.

No core measures for disability were recommended for evaluation by the EMSOP investigators and consultants. Collecting measurements on this parameter would be very challenging for several reasons. First, no one functional outcome measure has been validated for all the priority conditions. Second, when disability measurements are obtained they are usually obtained at intervals distal to the prehospital and emergency department service intervals. Third, disability measures are not routinely obtained on emergency patients, whether or not they are brought to the emergency department by EMS. Thus, there are no data sources available for linkage for the broad array of priority conditions. The fact that no core disability measure was recommended for evaluation does not imply that prehospital care researchers should not continue the search for an appropriate core disability measure. Even though such a measure may not be routinely collected on EMS patients, it may have great value for specific research projects. Later in this report, we discuss disability measures that may be applicable to specific priority conditions.

With regard to the outcome category discomfort, an expert panel identified relief of discomfort as : 1) the most relevant outcome parameter in both adult and pediatric priority conditions; 2) the prehospital intervention that may have the greatest impact on patients (Maio et al., 1999). We think that for the overwhelming majority of patients in the prehospital setting, the measurement of pain is the most appropriate measure of discomfort. Due to its importance, we have devoted the entire next section of the report to pain measurement.

Patient satisfaction is an outcome measure that has been used for a variety of purposes in health care (Allen and Rogers, 1996). Satisfaction has been identified as an indicator of quality care (Doering, 1983) and can also be used to assess the performance of health care

delivery at the organizational, unit and individual level. Patient satisfaction is also used in the development and evaluation of patient care models. Although we think that a measure of satisfaction should be obtained on all EMS patients, we were not able to recommend a satisfaction measure that should be evaluated in the prehospital care setting. We could find no report in the peer-reviewed literature of the development, evaluation and use of a patient satisfaction instrument, either generic or disease-specific, that has been used in the prehospital setting. Although there have been a number of studies that have addressed patient satisfaction in the emergency department, either as the primary or secondary outcome measure, only two instruments have reported information regarding their validity, reliability and limitations. It is readily apparent the field of prehospital care is in urgent need of an appropriately developed and tested instrument to assess patient satisfaction. One logical approach would be to develop a basic instrument that could be broadly applied, and eventually modified to address specific diseases. Researchers could begin developing prehospital patient satisfaction instruments by modifying existing measures such as the Group Health Association of America (GHAA) consumer satisfaction survey (Davies and Ware, 1991), or the Picker/Commonwealth survey (Pascoe, 1983). Furthermore, future research may be able to utilize the findings from the ongoing Consumer Assessment of Health Plan Study (CAHPS) sponsored by the Agency for Health Care Research and Quality (AHRQ).

The final outcome category, cost, is intended to capture not only the physical and personal outcomes that result from patient care, but also the costs and benefits of such care. The analyses of costs are typically studied using three models: 1) Cost-effectiveness analysis; 2) Cost-benefit analysis; and 3) Cost-utility analysis (Drummond et al., 1997). In a cost-effectiveness analysis, consequences are measured in natural units such as the amount of

disability or health care resources consumed. Typically, there is a single effect of interest, common to both alternatives, but achieved to different degrees. Using the cost-benefit analysis, consequences are measured in dollars. This is similar to cost-effectiveness analysis except that there may be single or multiple effects of interest that are not necessarily common to both alternatives and common effects may be achieved to different degrees by the alternatives. In an analysis of cost-utility, consequences are measured in quality adjusted life years. Typically, cost-utility analysis is conducted when effects are weighted by utility measures denoting the patient's preference for, or the overall desirability of, a particular outcome (Gold, et al., 1996). Further complicating the picture is whether only direct costs are considered (for example acute medical care) or indirect costs (days of work missed) and what the perspective is: patient, insurer, or society. Furthermore, when considering direct costs there is controversy with regard to whether to use charges to patients and/or insurers versus "true costs". Charges for similar service may vary based on the insurer and "true costs" for a various episode of care can be variable based on the measures used to determine those costs.

A number of studies have appeared in the peer-reviewed literature that have examined the issue of cost as an outcome of pre-hospital care. All of the studies are cost-effectiveness studies. Cost-benefit and cost-utility studies do not appear in the peer-reviewed literature to date. The topics of published studies include the cost-effectiveness of helicopter transport including an analysis of helicopter EMS for trauma patients (Gearhart, et al., 1997); a comparison of air medical compared to ground transport (Spaite, et al., 1994), and air medical helicopter crash survival enhancements (Dodd, 1994). Another topic of cost-effectiveness studies is the issue of out-of-hospital cardiac arrest (Valenzuela, et al., 1991). Similar studies have considered potential improvements to EMS for victims of out-of-hospital cardiac arrest

(Nichol, et al., 1996), and the cost-effectiveness of public access defibrillation (Nichol et al., 1997, 1998). Other topics appearing in the literature include expanding out-of-hospital care and its impact on hospital resource usage (Wade, 1996), the cost-effectiveness of trauma care (Elliott, 1996), and comparing ambulance dispatch protocols for nontraumatic abdominal pain (Lammers et al., 1995). The EMSOP investigators, however, could find no cost measure that could be uniformly applied to all EMS patients across many different systems. Future research is needed to develop, evaluate and implement cost determination methods that can be widely applied through the EMS community.

5.2.3 Pain

Inadequate pain control has also been recognized in the field of emergency medicine, both in the emergency department and in the prehospital setting. It has also been termed the "fifth vital sign". As discussed earlier, among priority conditions identified, discomfort had the highest weighted score for the top 3 adult first quartile conditions (minor trauma, respiratory distress and chest pain) and the first and third highest ranking children's first quartile conditions (minor trauma and respiratory distress). We think two measures of pain intensity should be evaluated for use among adults and older children in the prehospital setting. One measure uses an adjective response scale (ARS) [Jensen et al., 1986; McGuire, 1984; Keele, 1958; Ohnhaus and Adler, 1975; Woodforde and Merskey, 1972; Huskisson, 1974] and the other a numeric rating scale (NRS) [Jensen et al., 1986; Downie et al., 1978; Wilkie et al., 1990]. The first scale can be used to determine the quality of pain intensity and the latter to quantify pain intensity. Both of these scales have been found to be valid and reliable. These scales have also been found to be feasible to use in the emergency department.

Two prehospital studies suggest that these instruments may be feasible to use in the prehospital setting. The ARS could be used upon initial assessment by pre-hospital care personnel and prior to relinquishing care of the patient to the emergency department staff (Ricard-Hibon et al., 1997, 1999). The following classification is recommended: None, Slight, Moderate, Severe, Agonizing. If a patient reports any category other than “None” he/she is asked to give a numerical rating to their pain from 0-100, with 0 being no pain and 100 being the worst pain imaginable. The numeric scale should be used at initial assessment and prior to turning over care to the emergency department staff. For younger children, self-report of pain can be utilized using the Oucher Scale (Beyer et al., 1992; Beyer and Knott, 1998; Villarruel and Denyes, 1991). This scale has been found to be valid and reliable but has not been evaluated in the emergency department or prehospital setting. Pain assessment instruments for infants are complex to use and are not currently feasible for use in the prehospital setting (Grunau et al., 1990, 1998; Craig et al., 1973, 1984; Attia, et al., 1987; Barrier et al., 1989; Horgan and Choonara, 1996; Lawrence et al., 1993; Taddio et al., 1995).

5.2.4 Condition-Specific Risk Adjustment and Outcome Measures

Next, the EMSOP investigators identified risk adjustment measures and outcome measures that should be measured or collected for specific EMS conditions. Measures were identified by a systematic literature search and a structured review of articles pertaining to that measure (see Appendix III for a description of the process used to identify measures). In addition, determination of outcome measures included review of several authoritative texts. Measures were evaluated using the following attributes: time taken to complete, cost and training, scaling, reliability, feasibility of use in the prehospital setting, and potential use as an

outcome measure. After evaluation, the EMSOP investigators discussed each measure and a decision made to include or discard the measure. The methodology of the search and reviews, findings and recommended measures were presented to and reviewed by the consultant panel of experts.

5.2.4.1 *Major/minor trauma*

In phase one of the project, we found that 51% of pediatric transports and 36% of adult transports were for minor injury. Defining minor injury may be somewhat arbitrary. In the first portion of this project, the investigators used the Revised Trauma Score (RTS) to distinguish between major and minor trauma. An RTS of 11 or less was considered major trauma and an RTS of more than 11 was considered minor trauma (Champion et al., 1989). This is the criterion that has been recommended for use by field personnel to determine which patients need to go to a trauma center. Using this definition, 51% of pediatric transports and 36% of adult transports of the data obtained in phase one were classified as minor injury. This measure was selected because it is a standard measure in trauma care and evaluation and could be computed from the available data. However, this manner of defining “minor trauma” is a very crude one, in that it divides patients into those with higher and lower risk of death. Those falling into the low risk category based on RTS range from patients with sprained ankles to patients with splenic lacerations who are alert, oriented and have normal vital signs. Although a precise universal definition for minor trauma is lacking, it usually includes patients in whom the risk of death is minimal. While acknowledging the limitations of our method for defining minor and major trauma, we nonetheless will clarify which of the following recommended measures should be evaluated for use in minor trauma, major trauma

or both. For major and minor trauma all previously discussed core risk adjustment and outcome measures will be used. It is important to note that the core measure of discomfort (or pain), may be the outcome parameter that is most relevant to those with minor trauma.

5.2.4.1.1 Risk Adjustment Measures: Major/minor trauma

Physiologic measures for trauma can be measured in the prehospital setting and are affected by time and treatment. The first measure, the Revised Trauma Score (RTS) uses physiological parameters to quantify injury severity-Glasgow Coma Scale, systolic blood pressure and respiratory rate. Specific ranges of each parameter are assigned coded values and also weights. Unweighted coded values range from 0-12, with 12 being no significant physiologic derangement. A patient with an RTS of 11 or less is considered at risk for experiencing major traumatic injuries. The RTS has been found to be a valid measure of physiologic injury severity in children (Eichelberger et al., 1989).

The second measure, the Glasgow Coma Scale (GCS) is a physiological injury severity score and is a crude but effective measure of the global function loss before and after resuscitation and prior to definitive treatment. This measure has been previously described under core measures. Although the RTS and GCS are accepted physiologic measures to risk adjust among patients with major injury, there are no physiologic parameters that have been identified for risk adjusting among minor injury patients. Both these measures can be obtained in the prehospital interval.

The Abbreviated Injury Scale (AIS) is the most widely used and accepted measure of injury severity based on anatomic descriptors (AAAM, 1990). It is an ordinal scale ranging from 1 (minor injury) to 6 (maximum injury, non-survivable). For the 1990 version injuries

are grouped into 9 body regions. Several studies have confirmed the validity and utility of the AIS. The validity of AIS scoring for pediatric injury has been examined and several modifications made to accommodate scoring for children as well as adults. AIS scores should be determined for each patient. The AIS rates the severity of individual injuries only. The most widely used measure for combining AIS scores across body regions is the Injury Severity Score (ISS). This score was devised by Baker (1976) and was found to correlate better with mortality than did the AIS. The ISS is defined as the sum of squares of the highest AIS score in each of three most severely injured body regions. The highest score possible is 75. Low scores have been correlated with low probability of death and high scores with high probability of death. Although widely used, the ISS has been criticized for not taking into account the combined effect of multiple injuries within a single body region and underscoring the severity of isolated head injuries (Somers, 1981; Copes et al., 1988). The New Injury Severity Score (NISS) has been developed by Osler and Baker (1997) and addresses some of the limitations of the ISS while maintaining simplicity in definition and computation. The NISS is the sum of squares of the three most severe regions, regardless of body region. Initial studies suggest that the NISS offers a modest increase in predictive accuracy regarding mortality. One of the obstacles to determining the AIS and subsequently, the ISS and NISS is the need for complete and precise information on the extent of injury. This will require the sharing of both emergency department and inpatient information by receiving hospitals. Although the ISS score can be used for both major and minor injuries, there is very little variation in the ISS score among injury patients that are treated and released from the emergency department. Future research is needed to develop a valid and reliable minor severity score.

5.2.4.1.2 *Outcome Measures: Major/minor trauma*

Besides the usual core mortality measures, special mortality measures can be used in the evaluation of trauma care. With some of these methods, anatomical injury severity scores are combined with physiologic injury severity scores to calculate a probability for survival (Ps). One can then do a detailed examination of patients who died that were at or above a certain Ps, such as 0.75. Using TRISS or ASCOT methodology one can also compute various statistics that would allow one to compare the overall survival rate of one system, adjusted for Ps, with survival rates from other systems (Boyd et al., 1987; Markle et al., 1992). The preventable death rate can also be obtained through case review by a panel (Cales et al., 1984; Cayten et al., 1991; Maio et al., 1996; Esposito et al., 1995). Detailed information is collected only on those trauma patient who die. The method can be used even if injury severity scores of trauma patients are not routinely measured by the institutions caring for trauma patients within the EMS system. Using a well-defined structural review process will increase the reliability of the method (Maio et al., 1996).

The RTS and the GCS, which are used as risk adjusters, can also be used as an outcome measure to determine progression of the severity of the injury. These measures can be obtained in the prehospital interval or be obtained by linking to data sources from more distal intervals. This measure would be most useful among patients with major injury.

The Functional Independence Measure (FIM) is a measure of instrumental activities of daily living that is frequently used among patients admitted to trauma centers (Guyatt et al., 1996; Hamilton et al., 1987). It consists of 18 items that summarize level of independence in 6 areas: self-care, sphincter control, transfer mobility, ambulation, communication and social cognition. The FIM score is usually based on direct observation, but can also be derived using

a structured questionnaire completed by self-report. The questionnaire can be administered in person or over the phone. It has now been applied to measuring outcomes following spinal cord injury, brain injury and severe orthopedic trauma (Brenneman et al., 1997; Bergner et al., 1981; Heinemann et al., 1997; Hetherinton et al., 1995). A limitation of the FIM that needs to be considered is that it is not sensitive to variations in higher end functioning that would be more typical of many orthopedic and mild to moderate brain injuries. A pediatric version of the FIM (Wee FIM) has also been developed (McCabe et al., 1990). Within the last few years a measure of Functional Assessment Measure (FAM) was developed as an adjunct to the FIM and contains items related to motor, cognitive and psychological aspects of everyday functioning (Hall et al., 1993; Hall et al., 1994). Use of the FIM would require linkage to hospital data. The FIM is not a measure applicable to minor trauma patients.

The Medical Outcomes 36-Item Short Health Survey (SF-36) was designed as a generic health indicator of health status for use in large population based studies (Ware et al., 1992; Ware et al., 1993; Ware et al., 1994). The SF-36 consists of 35 items or questions that are scaled to measure eight health concepts: physical functioning, role limitations due to physical health problems, bodily pain, general health perceptions, vitality, social functioning, role limitations due to emotional problems, and general health. When using the SF-36 to examine post-injury outcome, some modification to questions about change in status or abilities is recommended to ascertain changes that may have occurred since the injury.

Two summary scores can be derived that measure physical and mental health respectively. The SF-36 can be self-administered or administered by an interviewer and takes only 5 to 10 minutes to complete. The SF-36 can also be incorporated as part of telephone interview. There are few published examples of its application to injury (Beaton et al., 1994;

Corrigan et al., 1998; Kopjar et al., 1996; Martin et al., 1997; McCabe, 1996). MacKenzie et al. (1998) has shown that SF-36 does not discriminate well among major trauma patients with and without head injury. However, by supplementing the SF-36 with 3 or 4 items, one can derive a subscore specific to cognitive function, that when used in combination with the standard eight subscores of the SF-36, provides a health profile more relevant to injury. The SF-36 has also been used to measure changes in functional outcome among patients with minor trauma (Kopjar, 1996). The SF-36 can definitely be used in the evaluation of major trauma and may also have a role in the evaluation of minor trauma. One of the problems with the SF-36 is that it is not routinely collected in all trauma patients. Therefore, it will most likely be a measure that can only be evaluated through specific research protocols that include collection of data that is not routinely obtained by the EMS system.

We think the FIM and SF-36 hold the most promise for measures of dysfunction among prehospital trauma patients. Currently, these are the two main functional outcome measures that are being used in a nationwide trauma evaluation funded by the CDC (National Study on Costs and Outcomes of Trauma Care (N-SCOT)). There are other measures of dysfunction that EMS researchers may also want to consider. The Sickness Impact Profile (SIP) consists of 136 statements about limitations in 12 categories: sleep and rest, emotional behaviors, body care and movement, eating, home management, mobility, social interaction, ambulation, alertness behavior, communication, recreation, and work. It is one of the most comprehensive measures of health status, addressing most dimensions of health with considerable depth (Bergner et al., 1981). It is also sensitive to a broad range of levels of dysfunction. The SIP, however, does take 20-25 minutes to administer, making it impractical for many applications. Shorter versions have been proposed for specific applications but

have not been widely validated (Gerety et al., 1994; Post et al., 1996; Sullivan et al., 1993). The SIP can be self-administered or administered by an interviewer. The SIP has been used in several studies describing the outcomes of spinal cord injury, head injury and orthopedic trauma (Beaton et al., 1994; Corrigan et al., 1998; Dikman et al., 1995; Fleming et al., 1998; Fuergemann et al., 1993; Gruen et al., 1995; Jurkovich et al., 1995; Richmond et al., 1998). In general, these studies support the validity of the SIP for trauma outcomes research. As with the FIM, the SIP would not be used in the evaluation of minor trauma. The Child Health Questionnaire (CHQ) was originally developed by Landgraf et al (1996) to measure the well-being of children and adolescents five years and older across 14 domains; a version of the CHQ applicable to toddlers is currently under development. Twelve of these domains can be combined into two summary measures of physical and psychosocial health status. Four different versions of the instrument exist consisting of 28 items, 50 items and 98 items respectively. It can be self-administered or interviewer administered and both parent and child versions exist for young and older children. To date, there are no published studies using this measure for injury. The Quality of Well-Being (QWB) Scale is a preference based measure of health that combines patient reported symptoms and disability into a single index that provides an expression of well being that ranges from zero for death to one for asymptomatic full functioning (Holbrook et al., 1998). It has been used to examine trauma outcomes, but still requires further evaluation.

Currently, there are no routine measures of satisfaction that are collected for trauma patients. Nor could we identify any measure that has been specifically evaluated in the trauma patient. As stated previously, we think that satisfaction measures should be applied

for every prehospital patient. Future research is required to develop and evaluate such a measure.

We could find no standardized method of cost determinations that could be applied to the broad range of prehospital care trauma patients. Various methods have been used to determine costs among trauma patients who are admitted to hospitals. Some hospitals may routinely collect some of these measures. However, no one measure or set of measures is recognized as the standard measure for cost. Future research will be needed to identify and/or develop and evaluate measures of cost relevant to prehospital trauma care.

5.2.4.2 Respiratory distress

In the first phase of this project we found that 13% of all adult transports and 10% of all pediatric transfers were for respiratory distress; these conditions ranked 2 and 3, respectively, among priority conditions. When considering recommended measures for the evaluation of respiratory distress in the prehospital setting, several considerations were taken into account. We considered that for the condition respiratory distress, a patient may have various underlying diseases such as asthma, congestive heart failure, COPD, and pneumonia. We also considered there were certain diseases specific to young children, to include croup and bronchiolitis. Prehospital care providers are trained, and prehospital protocols are developed to treat symptoms and signs, not specific disease categories. The very nature of emergent care often precludes the accurate diagnosis of the disease that is causing the symptoms. Therefore, we had to identify measures that could be used among several diseases. Another particularly challenging issue in the process was identifying measures that would be useful among a broad age range. Since the specific disease states that would fall

under the prehospital condition of respiratory distress have never been reported, we decided to identify measures that could be used for the specific diseases of asthma, COPD and congestive heart failure. The EMSOP investigators and consultants thought that these three conditions comprise the overwhelming majority of adult and pediatric transfers for respiratory distress. As with the condition of major/minor trauma, all core risk adjustment and outcome measures would be obtained for patients with respiratory distress. These core measures will not be discussed further in this section.

5.2.4.2.1 *Risk Adjustment Measures: Respiratory Distress*

Peak Expiratory Flow Rate (PEFR) is one of a group of forced expiratory flow measures taken at the point of total lung capacity or at the point of maximal inspiration (American Thoracic Society, 1987 and 1991). Wright first described the measurement of the PEFR, and the instrument used to obtain the measurement in 1959 (Wright, 1959). The PEFR is primarily an index of obstruction in large airways. The use of the PEFR to monitor patients with asthma has been widely adapted (Thoracic Society of New Zealand, 1996). Its use in the treatment and evaluation of chronic obstructive pulmonary disease has also been recommended (Aschraft et al., 1969; Tashkin, 1979). Studies in patients with congestive heart failure have found only mild decreases in PEFR in stable patients, but higher decreases in patients with severe symptoms of CHF (Hales et al., 1977; Light et al 1983, Peterman et al., 1987; Eriksson et al., 1987; Eichaker et al., 1988; Pison et al., 1989). Since Wright's first article, various other instruments have been developed to measure the PEFR. These have been inexpensive portable devices that have been calibrated using the Wright flow meter as a standard (Miller et al., 1992). A device that is frequently used today in the United States is

the mini-Wright peak flow meter (Wright, 1978). Children as young as four years of age can be taught to use portable devices to measure PEFr (Jones et al., 1990). The validity of the PEFr has been measured using the FEV₁ as a gold standard. In general, the PEFr correlates well with FEV₁ (Nowak et al., 1982; Vaughn et al., 1989). Others have obtained similar results in stable and asthmatic children and in normals. A study by Vaughn et al. (1989) noted correlations of 0.74. Studies conducted in emergency departments have found correlations ranging from 0.77-0.83 in asthma and 0.69 in COPD (Emmerman et al., 1989).

Oxygen saturation represents the percent saturation of available bound hemoglobin. The relationship between percent saturation and the partial pressure of oxygen in arterial blood (PaO₂) is described by the oxyhemoglobin dissociation curve. The pulse oximeter (SpO₂), is a fast growing and commonly utilized transcutaneous methodology that represents a combination of oximetry and plethysmographic technologies for screening, treatment planning and evaluation and research (Fait et al., 1985; Mihm et al., 1989; Yelderman et al., 1983). Evaluation of the accuracy and feasibility of pulse oximetry in the emergency department and prehospital setting have also been conducted (Jones et al., 1988; Aughey et al., 1991; McGuire et al., 1988). These studies have found the device accurate and its use very feasible. Precision of various pulse oximeters were noted to be fairly consistent at approximately 2% (Taylor et al., 1988). The anatomic site of choice for pulse ox probe placement is the finger (Webb et al., 1991). Limitations of this device include severe anemia and low-flow states (Webb et al., 1991).

Dyspnea can be defined as the unpleasant sensation of labored or difficult breathing and is synonymous with the term shortness of breath. This is main parameter of discomfort in respiratory distress (Gift et al., 1993). Factors related to the presence of dyspnea can be

classified as physiologic, psychologic, and situational or environmental (Gift et al., 1993).

The visual analogue dyspnea scale (VADS) is a self-reported measure that has shown to be a valid measure of dyspnea, as well as practical in the critical care setting (Gift, 1986; Gift et al., 1986; Aitken, 1969). The patient is asked to indicate the degree of shortness of breath experienced by marking the line at the level indicating his or her level of discomfort (Gift, 1989). One end represents “not at all breathless” and the other end “worst possible breathlessness”. Children must have fully developed communication skills in order to use these instruments. Studies in the non-emergency setting have reported that children as young as four could use the VADS.

The visual analogue scale has been validated in healthy volunteers as well as asthmatics and persons with COPD as well as in cancer patients suffering from a variety of underlying causes of dyspnea (Gift, 1989; Adams et al., 1985; Roberts et al., 1993; Mador et al., 1992; Subratty et al., 1994; Nosedá et al., 1992). The test-retest reliability of the VAS has been shown to be high (Muza, 1990). Aitken (1969) was the first to demonstrate that a visual analogue scale was a valid measure of dyspnea. He used a horizontally oriented scale. Gift, Plaut and Jacox (1986) expanded Aitken’s work by reorienting the scale to a vertical line and testing the scale in clinical situations using asthmatic patients and COPD. This reorientation facilitated ease and understanding of use by patients. The vertical VAS is a valid, reliable and easy-to-use instrument. It is an instrument that could be easily used in the prehospital setting (Gift, 1986).

5.2.4.2.2 Outcome Measures: Respiratory Distress

No special measures of mortality for respiratory distress were identified. The PEFR and Pulse Ox, risk adjustment measures for respiratory distress, can also be used as outcome measures for disease severity. These outcome measures can be obtained in the prehospital care interval or obtained through linkage of data from more distal service intervals. The VADS, used to initially measure discomfort for prehospital patients, can also be used as an outcome measure for discomfort. This outcome measure can be obtained in the prehospital care interval or, by linkage, from data of more distal service intervals.

No functional outcome measures that could be used for all patients with respiratory distress could be identified. Although there are disease specific measures of functional status that could be used among patients with respiratory distress (examples include the New York Heart Association Classification, St. George's Respiratory Questionnaire, Pulmonary Function Status Scale, Chronic Respiratory Disease Questionnaire). None of these measures are routinely collected on patients with respiratory distress who are treated and released from the emergency department or who are admitted to the hospital. Future research will be needed to determine the feasibility of using these and other specific respiratory disease functional status measures in conducting outcomes research and evaluation of EMS patients who present with respiratory distress.

Currently there are no routine measures of satisfaction that are routinely collected for patients with respiratory distress in the prehospital setting. Nor could we identify any measure that has been specifically evaluated among patients with respiratory distress. As stated in the section on core risk adjustment and outcome measures, we think that every

prehospital patient should have a measure of satisfaction obtained. Future research will need to develop and evaluate such a measure.

We could find no standardized method of cost determinations that could be applied to the broad range of prehospital care respiratory distress patients. Future research will be needed to identify, or develop, and evaluate measures of cost relevant to prehospital care of patients in respiratory distress.

5.2.4.3 *Cardiac Arrest*

Of all outcomes research undertaken in the out-of-hospital setting, the highest quality to date has been in the arena of non-traumatic cardiac arrest. The EMSOP investigators recommend the use of the Utstein Style (Cummins et al., 1991a). Utstein has already enjoyed wide acceptance and a large amount of work is underway using the guidelines for cardiac arrest research (AHA, 2000; Cummins et al., 1991b; Steill et al., 1998; Steill et al., 1999a, 1999b; Hsu et al., 1996; Grubb et al., 1996; Callicot et al., 1995; Cobbe et al., 1996; Nichol et al., 1999; The Brain Resuscitation Clinical Trial II Study Group, 1991; Cobbe et al., 1996).

Almost nothing has been done in the methodological development of cardiac arrest research in children. Thus, much work will have to be done before any cogent recommendations can be made for pediatric cardiac arrest outcomes research. There are rumors of the development of Utstein Style guidelines for pediatrics, but no accepted or authoritative methods are extant at present. Therefore, EMSOP cannot make evidence-based recommendations at this time.

5.2.4.4 Chest Pain

We were unable to identify any risk adjustment or outcome measures specific to chest pain. We recommend that for chest pain, the general advice provided in EMSOP IV, *Pain measurement in prehospital outcomes research* be followed. Future research should focus on appropriate and specific out-of-hospital risk adjustment and outcomes measures for chest pain.

5.2.4.5 Respiratory Arrest/Airway Obstruction

We were unable to identify any specific risk adjustment or outcome measures for these entities. At the very least we would recommend core risk adjustment and outcome measures for these patients. Future research will need to develop and evaluate appropriate condition specific risk adjustment and outcome measures for respiratory arrest/ airway obstruction.

5.2.4.6 Pediatric Seizures

We were unable to identify any specific risk adjustment or outcome measures for these entities. At the very least we would recommend core risk adjustment and outcome measures for these patients. Future research will need to develop and evaluate appropriate condition specific risk adjustment and outcome measures for pediatric seizures.

5.3 Finding 4: Identify stakeholders and constituencies important for EMS outcomes research.

A list of stakeholders was developed in the second year of the project and modified in the fourth year with the input of the steering committee and consultants (see Appendix IV). These stakeholders and constituencies represent physicians, prehospital care providers, EMS

and ambulance system administrators, government agencies, insurance companies and legislators. It is important that all project information disseminated be understandable to both the research community and EMS professionals at the state level and in the field.

A dialog was begun with the National Association of EMS Physicians in July, 1997 when the priority conditions were identified to alert leading prehospital researchers of the project and its progress. Since that time, annual updates have been provided to key EMS researchers at the Society for Academic Emergency Medicine Annual Meeting. The network of stakeholders and constituencies should continue to provide input in the development of the EMS outcomes research agenda, as well as serve as a resource for dissemination.

5.4 Finding 5: Develop a research dissemination plan.

The manuscripts developed as a result of this project will reach the desired audience and help pave the way for an EMS outcomes research agenda. Further information will be developed for the broad EMS research community, in dialog with the EMS office of NHTSA and using the list of stakeholders that has been developed by the project investigators and consultants. A one-day “think tank” was held on March 20, 2002 on Performance Measures in EMS that included a broad audience of the EMS community including state EMS directors, firefighters and others involved in prehospital care. Dr. Ron Maio presented EMSOP findings to ensure that the objectives of this project, particularly with regard to outcomes in prehospital care, were incorporated into the Performance Measures work group. On June 3, 2002, EMSOP results were presented at the National EMS Research Agenda Implementation Symposium sponsored by the National Association of EMS Physicians. Another goal of the symposium was to draft a plan for the implementation of the National EMS Research Agenda

recommendations. Through all of these efforts, the ultimate goal is to conduct EMS outcomes research that will improve the delivery of prehospital care, and will thus benefit the general public.

6.0 Limitations

One of the primary limitations of this project was the paucity of EMS databases. We had no nationally representative data sources which contained information regarding the frequency and nature of prehospital care in our country. The EMS Data Systems database was not a probability based nationally representative sample. However, we think it provides us with a reasonable estimate of the frequency and nature of EMS care in our country and do not think that use of a true probability based national sample would substantially change our project's findings. Another limitation was the lack of prehospital studies that addressed methodological issues pertaining to risk adjustment and outcome measurement of the priority conditions that were identified. Therefore, as stated in previous sections of this report, often the best we could do was make recommendations regarding the most promising measures that need to be evaluated for feasibility in the prehospital setting. However, we think we have provided an invaluable service to those wanting to evaluate prehospital care in that they now have some idea of what measures hold the greatest promise. Our study did not actually conduct any outcomes research. We cannot comment on what interventions for our priority conditions are effective. However, from the start, this was never the intention of the EMSOP project.

Our mission was to develop a "blueprint" and a "toolbox" for EMS outcomes research. Our project did not specifically address interfacility transport, air-medical

transport, and the treatment and non-transport of patients. However, we think that many of the measures we are recommending would be readily applicable to these situations and encourage researchers to evaluate the EMSOP recommended measures in these settings. Since our project focused on outcomes from acute prehospital emergency care we did not consider measures relevant to EMS injury/disease prevention activities. Nonetheless, we think effectiveness research regarding prevention services delivered by prehospital care providers is an important research area. We were unable to identify and recommend any condition-specific risk adjustment or Outcome measures for Respiratory Distress/Airway Obstruction and Pediatric Seizures. Prehospital researchers interested in these areas may need to conduct hospital-based evaluations to develop and validate appropriate condition specific measures that can then be evaluated in the prehospital setting.

7.0 Implications

One of the major implications of our study is that it underscores the importance of studying the effect of prehospital care on non-mortality outcomes, in particular, the relief of discomfort. In fact relief of discomfort was the outcome parameter EMS professionals identified as having the most potential impact for the majority of children and adults in the top quartile conditions. Although EMSOP investigators, consultants and national experts participating in EMSOP I support the concept of alleviating discomfort as an important prehospital intervention, it is not known to what extent this opinion exists among prehospital care providers or other stakeholders such as patients, professional associations representing health care providers, and state and federal agencies. It may be difficult for some to modify the current mission of prehospital care, "saving lives", into a new mission, "relieving

discomfort". While understanding the tenacity of traditional thinking, we do not think the importance of relief of discomfort in the prehospital arena should be any less important than it is in other health care arenas. It is important to investigate stakeholder attitudes toward prehospital management of discomfort, and the concomitant possibility for impeding prehospital relief of discomfort research or the application of its findings.

Another implication of our research is the importance of obtaining both core and condition specific risk adjustment measures and outcome measures from sources outside of the control of prehospital care systems. Data must be shared by health care facilities that participate in prehospital care systems. Current concerns regarding patient confidentiality will no doubt provide some challenges to this sharing of data. Nonetheless, failure to share such data will preclude any meaningful prehospital care outcomes research.

8.0 Future Research

The following are needed: 1) Studies to substantiate or refute the appropriateness of the measures suggested by the EMSOP investigators; 2) Studies to determine the feasibility and reliability of using outcome measures among all levels of prehospital care providers; 3) Studies to determine if feasibility and reliability vary based on condition and/or core risk adjustment measures (RAM). After initial research identifies feasible and reliable measurement instruments applicable to the prehospital setting, the next step is to use those instruments to describe the distribution of the values of that measure in the prehospital setting. Patients transported for minor trauma might benefit from prehospital pain control. Once the distribution of the different values of the measurements have has been described, the next step is to consider the relationship of various prehospital risk adjustment measurements to

outcome measures in the prehospital unit of service, as well as more distal outcome measures. After identifying these important relationships, the next step would be to develop interventions that effectively treat the priority conditions that have been identified. It is possible that studies to determine the relationship of prehospital risk adjustment measures to outcomes distal to the prehospital interval of service will occur in parallel with intervention studies. Future research must also address identifying appropriate measurements for the outcome parameters of satisfaction and cost. Appropriate risk adjustment and outcome measures also need to be identified and evaluated for infants and very young children.

9.0 Project Participants

The steering committee for the project was chaired by Ronald F. Maio, DO (University of Michigan), also the principal investigator. Other steering committee members included Herbert Garrison, MD (East Carolina University), and Daniel Spaitte, MD (University of Arizona), co-principal investigators. Other project staff from the University of Michigan on the steering committee included Jeffrey Desmond, MD, co-investigator; and Mary Ann Gregor, MHSA, project coordinator. Consultants on the project and their representative specialties included: Gene Cayten, M.D., Institute for Trauma and Emergency Care, New York Medical College, Valhalla, NY (Trauma and Emergency Surgery); John Chew, EMSSTAR Group, Annapolis, MD (EMS Policy); Ellen MacKenzie, Ph.D., School of Hygiene and Public Health, Johns Hopkins University, Baltimore, MD (Health Services Research); Ian Stiell, M.D., University of Ottawa, Ottawa, Ontario (EMS Research); Patricia O'Malley, M.D., Massachusetts General Hospital, and Emergency Medical Services for

Children, Massachusetts Department of Public Health, Boston, MA (Pediatric Emergency Medicine); and David Miller, M.B.A., Allina Health System, St. Paul, MN (Paramedicine).

10.0 List of Publications (see Appendix V)

Maio RF, Garrison HG, Spaite DW, Desmond JS, Gregor MA, Cayten CG, Chew JL, Hill EM, Joyce SM, MacKenzie EJ, Miller DR, O'Malley PJ, Stiell IG. Emergency Medical Services Outcomes Project I (EMSOP I): Prioritizing Conditions for Outcomes Research." *Annals of Emergency Medicine* 33:423-432,1999.

Spaite DW, Maio RF, Garrison HG, Desmond JS, Gregor MA, Stiell IG, Cayten CG, Chew JL, MacKenzie EJ, Miller DR, O'Malley PJ. Emergency Medical Services Outcomes Project II (EMSOP II): Developing the foundation and conceptual models for prehospital outcomes research. *Annals of Emergency Medicine* 37:657-663, 2001.

Garrison HG, Maio RF, Spaite DW, Desmond JS, Gregor MA, Stiell IG, Cayten CG, Chew JL, MacKenzie EJ, Miller DR, O'Malley PJ. Emergency Medical Services Outcomes Project III (EMSOP III): Risk adjustment measures for prehospital outcomes research. *Annals of Emergency Medicine* 40:79-88, 2002.

Maio RF, Garrison HG, Spaite DW, Desmond JS, Gregor MA, Stiell IG, Cayten CG, Chew JL, MacKenzie EJ, Miller DR, O'Malley PJ. Emergency Medical Services Outcomes Project IV (EMSOP IV): Pain measurement in prehospital outcomes research. *Annals of Emergency Medicine* 40:172-179, 2002.

Garrison HG, Maio RF, Spaite DW. Application of measurement tools to pediatric emergency medicine. *Ambulatory Pediatrics* 2:319-322, 2002.

Spaite DW, Maio RF, Garrison HG, Desmond JS, Gregor MA, Stiell IG, Cayten CG, Chew JL, MacKenzie EJ, Miller DR, O'Malley PJ. Emergency Medical Services Outcomes Project V(EMSOP V): Establishing the scope and methodological approach to prehospital outcomes and effectiveness research. *Draft*.

11.0 List of Presentations

What Works in Out -of -Hospital Care? Presented at Society of Academic Emergency Medicine's Annual Meeting, Washington, D.C., May 19-22, 1997.

Use of National Highway Traffic Safety Administration Uniform Out-of Hospital Data Elements in Statewide EMS Databases. Presented at Society of Academic Emergency Medicine's Annual Meeting, Washington, D.C., May 19-22, 1997.

The EMS Outcomes Project. Presented at the National Association of EMS Physicians, Mid-Year Scientific Assembly, Lake Tahoe, Nevada, July 9-11, 1997.

- Use of National Highway Traffic Safety Administration Uniform Out-of-Hospital Data Elements in Statewide EMS Databases. Presented at the National Association of EMS Physicians, Mid-Year Scientific Assembly, Lake Tahoe, Nevada, July 9-11, 1997.
- Invited presentation to National Highway Traffic and Safety Administration (NHTSA) Taskforce on EMS Education, "The EMS Outcomes Project", Alexandria, Virginia, June 29, 1998.
- Invited presentation to National Highway Traffic and Safety Administration (NHTSA) Taskforce on EMS Research Agenda, "The EMS Outcomes Project", Lake Tahoe, Nevada, July 13, 1998.
- Invited presentation at the National Association of State Emergency Medical Services Directors (NASEMSD) Annual Meeting, Ashville, North Carolina, October 29, 1998.
- Outcomes Research: Setting the Stage for the Future. Presented at the Society for Academic Emergency Medicine's 1999 Annual Meeting, Boston, Massachusetts, May 20-23, 1999.
- Application of Measurement Tools to Pediatric Emergency Medicine. Presented to the Ambulatory Pediatric Association and Emergency Medical Services for Children Program Conference: Improving Emergency Medical Services for Children Through Outcomes Research: An Interdisciplinary Approach, Reston, VA, March 30, 2001.
- Emergency Medical Services Outcomes Project (EMSOP): Application in Future Performance Measurement Systems. Presented at A Leadership Forum on Performance Measures in EMS, Arlington, VA, March 20, 2002.

References

- Adams L, Chronos N, Lane R, Guz A. The measurement of breathlessness induced in normal subjects: validity of two scaling techniques. *Clin Science* 69:7-16, 1985.
- Afschrift M, et al. Maximal expiratory and inspiratory flow in patients with chronic obstructive pulmonary disease: influence of bronchodilation. *Am Rev Respir Dis* 100:147-152, 1969.
- Aitken RCB. Measurement of feelings using visual analogue scales. *Proceedings of the Royal Society of Medicine* 62:989-993, 1969.
- Allen MM, Rogers WH. Consumer surveys of health plan performance: a comparison of content and approach and a look to the future. *The Joint Commission Journal on Quality Improvement* 22:775-794, 1996.
- American Heart Association. *Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care - International Consensus on Science*. Dallas: American Heart Association, 2000.
- American Thoracic Society. Standardization of Spirometry—1987 Update. *Am Rev Respir Dis* 136:1285-1298, 1987.
- American Thoracic Society. Lung Function Testing: selection of reference values and interpretive strategies. *Am Rev Respir Dis* 144:1202, 1991.
- Attia J, Amiel-Tison C, Mayer MN, et al. Measurement of postoperative pain and narcotic administration in infants using a new clinical scoring system. *Anesthesiology* 67:532, 1987.
- Aughey K, Hess D, Eitel D, Blecher K, Cooley M, Ogden C, Sabulsky N. An evaluation of pulse oximetry in prehospital care. *Ann Emerg Med* 20:887-891, 1991.
- Baker SP, O'Neill B. The Injury Severity Score: an update. *J Trauma* 16:882-5, 1976.
- Barrier G, Attia J, Mayer MN, et al. Measurement of post-operative pain and narcotic administration in infants using a new clinical scoring system. *Intensive Care Med* 15(suppl):37-39, 1989.
- Beaton DE, Bombardier C, Hogg-Johnson S. Choose your tool: a comparison of the psychometric properties of five generic health status instruments in workers with soft tissue injuries. *Qual Life Res* 3:50-56, 1994.
- Bergner M, Bobbitt RA, Carter WB, et al. The SIP: development and final revision of a health status measure. *Med Care* 19:787, 1981.

- Beyer JE, Denyes MJ, Villaruel AM. The creation, validation and continuing development of the Oucher: a measure of pain intensity in children. *J Pediatr Nurs* 7:335, 1992.
- Beyer JE, Knott CB. Construct validity estimation for the African-American and Hispanic versions of the Oucher Scale. *J Pediatr Nurs* 13:20, 1998.
- The Brain Resuscitation Clinical Trial II Study Group. A randomized clinical trial of calcium entry blocker administration to comatose survivors of cardiac arrest: design, methods, and patient characteristics. *Controlled Clin Trials* 12:525-545, 1991.
- Brazier J, Usherwood T, Harper R, Thomas K. Deriving a preference-based single index from the UK SF-36 Health Survey. *J Clin Epidemiol* 51:1115-1128, 1998.
- Brenneman FD, Katyal D, Boulanger BR, et al. Long-term outcomes in open pelvic fractures. *J Trauma* 42:773-777, 1997.
- Brice JH, Garrison HG, Evans AT. Study design and outcomes in out-of-hospital research: a ten-year analysis. *Acad Emerg Med* 3:422, 1996.
- Cales RH. Trauma mortality in Orange County: the effect of a regional trauma system. *Ann Emerg Med* 13:1-10, 1984.
- Callahan M. Quantifying the scanty science of prehospital emergency care. *Ann Emerg Med* 30:785-790, 1997.
- Callicot C, Valenzuela TD, Spaite DW, Clark LL. Survival and quality of life after cardiac arrest. *Acad Emerg Med* 2:433, 1995.
- Cayten CG, Stahl WM, Agarwal N, Murphy JG. Analyses of preventable deaths by mechanism of injury among 13,500 trauma admissions. *Ann Surg* 214:510-520, 1991.
- Champion HR, Sacco WJ, Copes WS, Gann DS, Gennarelli TA, Flanagan ME. A revision of the Trauma Score. *J Trauma* 29:623-629, 1989.
- Cobbe SM, Dalziel K, Ford I, Marsden AK. Survival of 1476 patients initially resuscitated from out of hospital cardiac arrest. *BMJ* 312:1633-1637, 1996.
- Copes WS, Champion HR, Sacco WJ, Lawnick MM, Keast SL, Bain LW. The injury severity score revisited. *J Trauma* 28:69-77, 1988.
- Corrigan JD, Smith-Knapp K, Granger CV. Outcomes in the first 5 years after traumatic brain injury. *Arch Phys Med Rehabil* 79:298-305, 1998.
- Craig KD, Grunau RVE. Neonatal pain perception and behavioural measurement. In: Anand KJS, McGrath PJ. *Pain in neonates*. Amsterdam: Elsevier, 1973.

- Craig KD, McMahan RJ, Morison JD, et al. Developmental changes in infant pain expression during immunization injections. *Soc Sci Med* 19:1331, 1984.
- Cummins RO, Chamberlain DA, Abramson NS, et al: Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: the Utstein style. *Circulation* 84:960-975, 1991a.
- Cummins RO, Ornato JP, Thies WH, Pepe PE. Improving survival from sudden cardiac arrest: the "chain of survival" concept. *Circulation* 83:1832-1847, 1991b.
- Davies AR, Ware JE. *GHAA's consumer satisfaction survey and user's manual*. Washington, DC: Group Health Association of America, 1991.
- Delbridge TR, Bailey B, Chew JL, et al. EMS agenda for the future: where we are... where we want to be. *Ann Emerg Med* 31:251-263, 1998.
- Dikman SS, Ross BL, Machamer JE, Temkin NR. One year psychosocial outcome in head injury. *J International Neuropsych Soc* 1:67-77, 1995.
- Dodd RS. The cost-effectiveness of air medical helicopter crash survival enhancements. *Air Medical Journal* 13:281-93, 1994.
- Doering, ER. Factors influencing inpatient satisfaction with care. *Quality Review Bulletin* 9:291-299, 1983.
- Donabedian A. Advances in health assessment conference discussion panel. *J Chronic Dis* 40(suppl1):183S-191S, 1987.
- Downie WW, Leatham PA, Rhind VM. Studies with pain rating scales. *Ann Rheum Dis* 37:378-381, 1978.
- Drummond MF, O'Brien B, Stoddart GL. *Methods for the economic evaluation of health care programmes*. NY: Oxford University Press, 1997.
- Eichacker PQ, Seidelman MJ, Rothstein MS, Lejerntel T. Methacholine bronchial reactivity testing in patients with chronic congestive heart failure. *Chest* 93:336-338, 1988.
- Eichelberger MR, Bowman LM, Sacco WJ, Mangubat EA, Lowenstein AD, Gotschall CS. Trauma score versus revised trauma score in TRISS to predict outcome in children with blunt trauma. *Ann Emerg Med* 18:939-42, 1989.
- Elliott DC, Rodriguez A. Cost effectiveness in trauma care. *Surg Clin North Am* 76:47-62, 1996.

- Emerman CL, Cydulka RK. Use of peak expiratory flow rate in emergency department evaluation of acute exacerbation of chronic obstructive pulmonary disease. *Ann Emerg Med* 27:159-163, 1996.
- Eriksson H, Caidahl K, Larsson B, Ohlson LO, Welin L, Wilhelmsen L. Cardiac and pulmonary causes of dyspnea—validation of a scoring test for clinical-epidemiological use: the study of men born in 1913. *Eur Heart J* 8:1007-1014, 1987.
- Esposito TJ, Sanddal ND, Hansen JD, et al. Analysis of preventable trauma deaths and inappropriate trauma care in a rural state. *J Trauma* 39:955, 1995.
- Fagerman C, Frandsen PA, Rock ND. Residual impairment after lower extremity fracture. *J Trauma* 45:123-126, 1998.
- Fait CD, Wetzell RC, Dean JM, et al. Pulse oximetry in critically ill children. *J Clin Monit* 1:232-235, 1985.
- Fleming JM, Strong J, Ashton R. Cluster Analysis of self-awareness levels in adults with traumatic brain injury and relationship to outcome. *J Head Trauma Rehab* 13:39-51, 1998.
- Fletcher RH, Fletcher SW, Wagner EH. *Clinical Epidemiology: The Essentials*. 2nd edition. Baltimore, Maryland: Williams and Wilkins; 1988.
- Garrison HG, Maio RF, Spaite DW, Desmond JS, Gregor MA, Stiell IG, Cayten CG, Chew JL, MacKenzie EJ, Miller DR, O'Malley PJ. Emergency Medical Services Outcomes Project III (EMSOP III): the role of risk adjustment in out-of-hospital outcomes research. *Ann Emerg Med* 40:79-88, 2002.
- Gearhart PA, Wuerz R, Localio AR. Cost-effectiveness analysis of helicopter EMS for trauma patients. *Ann Emerg Med* 30:500-6, 1997.
- Gerety MB, Cornell, JE, Mulrow CD, et al. The Sickness Impact Profile for Nursing Homes (SIP-NH). *J Gerontol* 49:M2-M8, 1994.
- Gift AG. Validation of a vertical visual analogue scale as a measure of clinical dyspnea. *Am Rev Respir Dis* 133:A163, 1986.
- Gift AG. A Dyspnea assessment guide. *Crit Care Nurse* 9:79-86, 1989.
- Gift AG, Plaut SM, Jacox AK. Psychologic and physiologic factors related to dyspnea in chronic obstructive pulmonary disease. *Heart Lung* 15:595-602, 1986.
- Gift AG, Pugh LC. Dyspnea and fatigue. *Adv Clin Nurs Res* 28:373-384, 1993.

- Gold MR, Siegel JE, Russell LB, et al. *Cost-effectiveness in health and medicine*. NY: Oxford University Press, 1996.
- Grubb NR, O'Carroll R, Cobbe SM, Sirel J, Fox KA. Chronic memory impairment after cardiac arrest outside hospital. *Br Med J* 313:143-146, 1996.
- Gruen GS, Leit ME, Gruen RJ, et al. Functional outcome of patients with unstable pelvic ring fractures stabilized with open reduction and internal fixation. *J Trauma* 39:838-844, 1995.
- Grunau RVE, Johnston CC, Craig KD. Neonatal facial and cry responses to invasive and non-invasive procedures. *Pain* 42:295, 1990.
- Grunau RVE, Oberlander TF, Holsti L, et al. Bedside application of the neonatal facial coding system in pain assessment of premature neonates. *Pain* 76:277, 1998.
- Guyatt GH, Jaeschke R, Feeny DH, Patrick DL. In: Measurements in clinical trials: choosing the right approach. *Quality of life and pharmacoeconomics in clinical trials*. Spilker B (ed) Philadelphia, PA: Lippincott-Raven Publishers, 1996.
- Hales CA, Kazemi H. Pulmonary function after acute uncomplicated myocardial infarction. *Chest* 72:350-358, 1977.
- Hall KM, Hamilton BB, Gordan WA, Zasler ND. Characteristics and comparisons of functional assessment indices: Disability Rating Scale, Functional Independence Measure, and Functional Assessment Measure. *J Head Trauma Rehabil* 8:60-74, 1993.
- Hamilton BB, Granger CV, Sherwin FS, et al. A uniform national data system for medical rehabilitation. In: Fuhrer MJ (ed). *Rehabilitation outcomes: analysis and measurement*. Baltimore, MD: Paula H. Brookes, 1987.
- Heinemann AW, Kirk P, Hastie BA, et al. Relationships between disability measures and nursing effort during rehabilitation for patients with traumatic brain injury and spinal cord injury. *Arc Phys Med Rehabil* 78:143-149, 1997.
- Hetherington H, Earlam RJ, Kirk CJ. The disability status of injured patients measured by the Functional Independence measure (FIM) and their use of rehabilitation services. *Injury* 26:97-101, 1995.
- Holbrook TL, Anderson JP, Sieber WJ, et al. Outcome after major trauma: discharge and 6-month follow-up results from the Trauma Recovery Project. *J Trauma* 45:315-324, 1998.
- Horgan M, Choonara I. Measuring pain in neonates: an objective score. *J Pediatr Nurs* 8:24, 1996.

- Hsu JW, Madsen CD, Callaham ML. Quality-of-life and formal functional testing of survivors of out-of-hospital cardiac arrest correlates poorly with traditional neurologic outcome scales. *Ann Emerg Med* 28:597-605, 1996.
- Iezzoni LI. Using risk-adjusted outcomes to assess clinical practice: an overview of issues pertaining to risk adjustment. *Ann Thorac Surg* 58:1822-1826, 1994.
- Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison of methods. *Pain* 27:117-126, 1986.
- Jones J, Heiselman D, Cannon L, Gradisek R. Continuous emergency department monitoring of arterial saturation in adult patients with respiratory distress. *Ann Emerg Med* 17:463-468, 1988.
- Jones KP, Mullee MA. Measuring peak expiratory flow in general practice: comparison of mini Wright peak flow meter and turbine spirometer. *Br Med J* 300:1629-1631, 1990.
- Jurkovich GJ, Mock C, MacKenzie EJ, et al. The Sickness Impact Profile as a tool to evaluate functional outcome in trauma patients. *J Trauma* 39:625-631, 1995.
- Kaufmann CR, Rivara FP, Maier RV. Pediatric trauma: need for surgical management. *J Trauma* 29:1120-5, 1989.
- Keele KD. The pain chart. *Lancet* 2:6-8, 1958.
- Koenig KL. Quo Vadis: "Scoop and run", "stay and play", or "treat and street"? *Acad Emerg Med* 2:477-479, 1995.
- Koenig KL. Unscheduled access to health care: reengineering the 911 system. *Acad Emerg Med* 3:986-991, 1996.
- Kopjar B. The SF-36 health survey: a valid measure of changes in health status after injury. *Inj Prev* 2:135-9, 1996.
- Lammers RL, Roth BA, Utecht T. Comparison of ambulance dispatch protocols for nontraumatic abdominal pain. *Ann Emerg Med* 26:579-89, 1995.
- Lawrence J, Alcock D, McGrath P, et al. The development of a tool to assess neonatal pain. *Neonatal Netw* 12:59, 1993.
- Lindgraf JM, Abetz LN. Measuring health outcomes in pediatric populations: issues in psychometrics and application. In: *Quality of Life and Pharmacoeconomics in Clinical Trials*. Spilker B (ed.), Philadelphia: Lippincott-Raven Publishers, 1996.

- Light RW, George RB. Serial pulmonary function in patients with acute heart failure. *Arch Intern Med* 143:429-433, 1983.
- MacKenzie EJ, Morris JA, Jurkovich GJ, et al. Return to work following injury: the role of economic, social, and job-related factors. *Am J Public Health* 88:1630-7, 1998.
- Mador MJ, Kufel TJ. Reproducibility of visual analog scale measurements of dyspnea in patients with chronic obstructive pulmonary disease. *Am Rev of Respir Distress* 146:82-87, 1992.
- Maio RF, Burney RE, Gregor MA, Baranski MG. A study of preventable trauma mortality in rural Michigan. *J Trauma* 41:83-90, 1996.
- Maio R, Garrison HG, Spaite DW, Desmond JS, Gregor MA, Cayten CG, Chew JL, Hill EM, Joyce SJ, Mackenzie EJ, Miller DR, O'Malley PJ, Stiell IG. Emergency Medical Services Outcomes Project I (EMSOP I): prioritizing conditions for outcomes research. *Ann Emerg Med* 33:423-432, 1999.
- Maio RF, Garrison HG, Spaite DW, Desmond JS, Gregor MA, Stiell IG, Cayten CG, Chew JL, MacKenzie EJ, Miller DR, O'Malley PJ. Emergency Medical Services Outcomes Project IV (EMSOP IV): pain measurement in out-of-hospital outcomes research. *Ann Emerg Med* 40:172-179, 2002.
- Markle J, Cayten CG, Byrne DW, Moy F, Murphy JG. Comparison between TRISS and ascot methods in controlling for injury severity. *J Trauma* 33:326-332, 1992.
- Martin DP, Engelberg R, Agel J, Swiontkowski MF. Comparison of the Musculoskeletal Function assessment questionnaire with the Short Form-36, the Western Ontario and McMaster Universities Osteoarthritis Index, and the Sickness Impact Profile Health-Status Measures. *J Bone Joint Surg [Am]* 79(9):1323-35, 1997.
- McCabe MA, Granger CV. Content validity of a pediatric Functional Independence Measure. *Appl Nurs Res* 3:120-122, 1990.
- McCabe MA. Pediatric Functional Independence Measure: clinical trials with disabled and nondisabled children. *Appl Nurs Res* 9:136-138, 1996.
- McGuire DB. The measurement of clinical pain. *Nurs Res* 33:152-156, 1984.
- McGuire TJ, Pointer JE. Evaluation of a pulse oximeter in the prehospital setting. *Ann Emerg Med* 17:1058-1062, 1988.
- Mihm FG, Halperin BD. Noninvasive detection of profound arterial desaturations using a pulse oximetry device. *Anesthesiology* 62:85-87, 1985.

- Muza SR. Comparison of scales used to quantitate the sense of effort to breathe in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 141:909-913, 1990.
- National Highway Traffic Safety Administration. *EMS outcomes evaluation: key issues and future directions. Proceedings from the NHTSA workshop on methodologies for measuring morbidity outcomes in EMS*. Washington, DC: U.S. DOT; April 11-12, 1994.
- National Highway Traffic Safety Administration. *Emergency medical services: agenda for the future* (DOT HS 808 441). Washington, DC: U.S. DOT, August 1996.
- National Highway Traffic Safety Administration. *Uniform pre-hospital emergency medical services data conference*. Washington, DC: U.S. DOT; August 16-18, 1993.
- Nichol G, Stiell IG, Hebert P, Wells GA, Vandemheen K, Laupacis A. What is the quality of life for survivors of cardiac arrest? A prospective study. *Acad Emerg Med* 6:95-102, 1999.
- Nichol G, Hallstrom A, Ornato JP, et al. Potential cost-effectiveness of public access defibrillation in the United States. *Circulation* 97:1315-20, 1998.
- Nichol G, Laupacis A, Stiell IG, et al. Cost-effectiveness analysis of potential improvements to emergency medical services for victims of out-of-hospital cardiac arrest. *Ann Emerg Med* 27:711-20, 1996.
- Nichol G, Ornato J, Riegel BJ, et al. Potential cost-effectiveness of public access defibrillation in the United States: a decision analysis. *Circulation* 96(8S):183-I, 1997.
- Nosedá A, Schmerber J, Prigogine T, Yernault JC. Perceived effect on shortness of breath of an acute inhalation of saline or terbutaline: variability and sensitivity of a visual analogue scale in patients with asthma or COPD. *Eur Respir J* 5:1043-1053, 1992.
- Nowak RM, Pensler MI, Sarkar DD, et al. Comparison of peak expiratory flow and FEV₁ admission criteria for acute bronchial asthma. *Ann Emerg Med* 11:64-69, 1982.
- Offner PJ, Jurkovich GJ, Gurney J, Rivara FP. Revision of TRISS for intubated patients. *J Trauma* 32:32-35, 1992.
- Ohnhaus EE, Adler R. Methodological rating scale and the visual analogue scale. *Pain* 1975;1:379-384
- Osler T, Baker SP, Long W. A modification of the injury severity score that both improves accuracy and simplifies scoring. *J Trauma*. 43:922-925.
- Pascoe GC. Patient satisfaction in primary health care: a literature review and analysis. *Evaluation and Program Planning* 6:185-210, 1983.

- Pearson GA, Stickley J, Shann F. Calibration of the paediatric index of mortality in UK paediatric intensive care units. *Arch Dis Child* 84:125-128, 2001.
- Peterman W, Barth J, Entzian P. Heart failure and airway obstruction. *Int J Cardiol* 17:207-209, 1987.
- Pison C, Malo JL, Rouleau JL, Chalaoui J, Ghezzi H, Malo J. Bronchial hyperresponsiveness to inhaled methacholine in subjects with chronic left heart failure at a time of exacerbation and after increased diuretic therapy. *Chest* 96:230-235, 1989.
- Post MW, de Bruin A, de Witte L, Schrijvers A. The SIP68: a measure of health-related functional status in rehabilitation medicine. *Arch Phys Med and Rehabil* 77:440-445, 1996.
- Relman AS. Assessment and accountability: the third revolution in medical care. *N Engl J Med* 319:1220-1222, 1988.
- Raskin IE, Maklan CW. Medical treatment effectiveness research: a view from inside the Agency for Health Care Policy and Research. *Eval Health Prof* 14:161-186, 1991.
- Reines HD, Bartlett RL, Chudy NE, et al. Is advanced life support appropriate for victims of motor vehicle accidents: the South Carolina highway trauma project. *J Trauma* 28:563-569, 1988.
- Ricard-Hibon A, Chollet C, Saada S, et al. A quality control program for acute pain management in out-of-hospital critical care medicine. *Ann Emerg Med* 34:738-744, 1999.
- Ricard-Hibon, Leroy N, Magne M, et al. Evaluation of acute pain in prehospital emergency medicine. *Ann Fr Anesth Reanim* 16:945-949, 1997.
- Richmond TS, Kauder D, Schwab CW. A prospective study of predictors of disability at 3 months after non-central nervous system trauma. *J Trauma* 44:635-642, 1998.
- Roberts DK, Thorne SE, Pearson C. The experience of dyspnea in late-stage cancer, patients' and nurses' perspectives. *Cancer Nursing* 16:310-320, 1993.
- Roper WL, Winkenwerder W, Hackbarth GM, et al. Effectiveness in health care: an initiative to evaluate and improve medical practice. *N Engl J Med* 319:1197-1202, 1988.
- Seidel JS, Henderson D, Tittle S, et al. Priorities for research in emergency medical services for children: results of a consensus conference. *Pediatr Emerg Care* 15:55-58, 1999.
- Shann F, Pearson G, Slater A, Wilkinson K. Paediatric index of mortality (PIM): a mortality

- prediction model for children in intensive care. *Intensive Care Med* 23:201-207, 1997.
- Smith JP, Bodai BI, Hill AS, et al. Prehospital stabilization of critically injured patients: a failed concept. *J Trauma* 25:65-70, 1985.
- Somers RL. *The probability of death score: an improvement of the ISS*. Laboratory for Public Health and Economics, Odense University Hospital. Odense, Denmark, 1982.
- Spaite DW. Outcomes analysis in EMS systems (editorial). *Ann Emerg Med* 22:1310-1311, 1993.
- Spaite DW, Criss EA, Valenzuela TD. Air medical transport of trauma patients: air versus ground transport. *Top Emerg Med* 16(4):73-80, 1994.
- Spaite DW, Criss EA, Valenzuela TD, et al. Developing a foundation for the evaluation of expanded-scope EMS: a window of opportunity that cannot be ignored. *Ann Emerg Med* 30:791-796, 1997.
- Spaite DW, Criss EA, Valenzuela TD, et al. Emergency medical services systems research: problems of the past, challenges of the future. *Ann Emerg Med* 26:146-152, 1995.
- Spaite DW, Maio R, Garrison HG, Desmond JS, Gregor MA, Stiell IG, Cayten CG, Chew JL, Mackenzie EJ, Miller DR, O'Malley PJ. Emergency Medical Services Outcomes Project II (EMSOP II): Developing the foundation and conceptual models for out-of-hospital outcomes research. *Ann Emerg Med* 37:657-663, 2001.
- Stiell IG, Spaite DW, Wells GA, Lyver MB, Munkley DP, Field BJ, Dagnone E, Maloney JP, Jones GR, Luinstra LG, Jermyn BD, Ward R, De Maio VJ. Modifiable factors associated with improved cardiac arrest survival in a multicenter basic life support/defibrillation system: OPALS Study Phase I Results. *Ann Emerg Med* 33:44-50, 1999a.
- Stiell IG, Wells GA, Field BJ, Spaite DW, De Maio VJ, Ward RE, Munkley DP, Lyver MB, Luinstra LG, Campeau T, Maloney JP, Dagnone E, for the OPALS Study Group. Improved out-of-hospital cardiac arrest survival through the inexpensive optimization of an existing defibrillation program, OPALS Study Phase II. *JAMA* 281:1175-1181, 1999b.
- Subratty AH, Manraj M, Baligadoo S. A new visual analogue scale for assessment of dyspnoea in congestive heart failure *Int J Clin Pharm Ther* 32:259-260, 1994.
- Sullivan M, Ahlmen, Bjelle A, et al. Health status assessment in rheumatoid arthritis II. Evaluation of a modified Shorter Sickness Impact Profile. *J Rheumatol* 20:1500-1507, 1993.
- Taddio A, Nulman I, Koren BS, et al. A revised measure of acute pain in infants. *J Pain Symptom Manage* 10:456, 1995.

- Tashkin DP. The UCLA population studies of chronic obstructive respiratory disease: II. Determination of reliability and estimation of sensitivity and specificity. *Environ Res* 20:403-424, 1979.
- Taylor MB, Whitwam JG. The accuracy of pulse oximeters. *Anaesthesia* 43:229-232, 1988.
- Thoracic Society of New Zealand. Peak flow meter use in asthma management. *Med J Australia* 164:727-730, 1996.
- Valenzuela TD, Criss EA, Spaite, et al. Cost-effectiveness analysis of paramedic emergency medical services in the treatment of prehospital cardiopulmonary arrest. *Ann Emerg Med* 20:944-5, 1991.
- Vaughan TR, Weber RW, Tipton WR, et al. Comparison of PEFR and FEV₁ in patients with varying degrees of airway obstruction. *Chest* 95:558-562, 1989.
- Villarruel AM, Denyes MJ. Pain assessment in children: theoretical and empirical validity. *Ans Adv Nur Sci* 14:32, 1991.
- Wade P. Evolving prehospital research; health service costing: a new perspective. *Australian Journal of Emergency Care* 3:32, 1996.
- Ware JE, Kosinski M, Keller SD. *SF-36 physical and mental health summary scores: a user's manual*. Boston, MA: The Health Institute, New England Medical Center, 1994.
- Ware JE, Sherbourne CD. The MOS 36-item Short-Form Health Survey (SF-36) I. Conceptual framework and item selection. *Med Care* 30:473-483, 1992.
- Ware JE, Snow KK, Kosinski M, et al. *SF-36 Health Survey: manual and interpretation guide*. Boston, MA: The Health Institute, New England Medical Center, 1993.
- Webb RK, Ralston AC, Runciman WB. Potential errors in pulse oximetry. *Anaesthesia* 46:207-212, 1991.
- Wilkie D, Lovejoy N, Dodd M; et al. Cancer pain intensity measurement: concurrent validity of three tools—finger dynameter, pain intensity number scale, visual analogue scale. *Hosp J* 6:1-13, 1990.
- Woodforde JM, Merskey H. Some relationships between subjective measures of pain. *J Psychosom Res* 16:173-178, 1972.
- Wright BM. A miniature Wright peak-flow meter. *Br Med J* ii:1627-1628, 1979.

Wright BM, McKerrow CB. Maximum forced expiratory flow rate as a measure of ventilatory capacity with the description of a new portable instrument for measuring it. *Br Med J* 2:1041-1047, 1959.

Yelderman M, New N. Evaluation of pulse oximetry. *Anesthesiology* 59:349-352, 1983.

Zuckerman GB, Gregory PM, Santos-Damiani SM. Predictors of death and neurologic impairment in pediatric submersion injuries. The Pediatric Risk of Mortality Score. *Arch Pediatr Adolesc Med* 152:134-140, 1998.

Appendix I

Questionnaire respondents and their affiliations

Questionnaire respondents and their affiliations

Bob Bailey, MA, North Carolina Office of EMS, Raleigh, NC

Ron G. Benoit, BS, EMT-P, University of Arizona, Tucson, AZ

Nicholas Benson, MD, East Carolina University, Greenville, NC

Dale J. Berry, EMT-P, Huron Valley Ambulance, Ann Arbor, MI

Richard A. Bissell, PhD, University of Maryland, Baltimore County, Baltimore, MD

Marni J. Bonnin, MD, Medical Center East Hospital, Birmingham, AL

William E. Brown Jr., RN, MS, NREMT-P, National Registry of EMTs, Columbus, OH

Lawrence H. Brown, EMT-P, SUNY Health Sciences Center, Syracuse, NY

C. Gene Cayten MD, MPH, New York Medical College, Valhalla, NY

John L. Chew Jr., MS, The EMSSTAR Group, Annapolis, MD

David C. Cone, MD, Allegheny University of the Health Sciences, Pittsburgh, PA

Drew E. Dawson, EMT-P, Montana EMS Bureau, Helena, MT

Theodore R. Delbridge, MD, MPH, University of Pittsburgh, Pittsburgh, PA

Jeffrey S. Desmond, MD, University of Michigan, Ann Arbor, MI

Herbert G. Garrison, MD, MPH, East Carolina University, Greenville, NC

Michael R. Gunderson, EMT-P, Institute for Prehospital Medicine, Tempe, AZ

Keith Holtermann, RN, MBA, MPH, George Washington University, Washington, D.C.

Richard C. Hunt, MD, East Carolina University, Greenville, NC

B. Tilman Jolly, MD, George Washington University, Washington, DC

Jon R. Krohmer, MD, Kent County EMS, Grand Rapids, MI

Ronald F. Maio, DO, MS, University of Michigan, Ann Arbor, MI

Questionnaire respondents and their affiliations (continued)

Dan Manz, BS, EMT-I, Vermont EMS Division, Burlington, VT

Steven A. Meador, MD, MPH, Hershey Medical Center, Hershey, PA

David R. Miller, BA, Health Span Transportation Services, St. Paul, MN

Keith Neely, MPA, EMT-P, Oregon Health Sciences University, Portland, OR

Patricia J. O'Malley, MD, Massachusetts General Hospital, Boston, MA

W. Taylor Payson, BA, MBA, EMT-P, Critical Air Medicine, Tucson, AZ

Ronald G. Pirrallo, MD, MHSA, Medical College of Wisconsin, Milwaukee, WI

Michael R. Sayre, MD, University of Cincinnati, Cincinnati, OH

Dena L. Smith, EMT-P, MSU/Kalamazoo Center for Medical Studies, Kalamazoo, MI

Daniel W. Spaite, MD, University of Arizona, Tucson, AZ

Ian G. Stiell, MD, MSc, Ottawa University, Ottawa, Ontario

Robert A. Swor, DO, William Beaumont Hospital, Royal Oak, MI

Vincent P. Verdile, MD, Albany Medical College, Albany, NY

Roger White, MD, Mayo Clinic, Rochester, MN

Richard Wuerz, MD, Brigham & Women's Hospital, Boston, MA

Michael Yee, EMT-P, University of Pittsburgh, Pittsburgh, PA

Appendix II

Sample item from questionnaire

Sample item from questionnaire

For this condition, how would you rate the potential impact of the following six outcome categories for patients < 15?

		Impaired	Limit	Alleviate		Cost
<u>Condition</u>	<u>Survival</u>	<u>Physiology</u>	<u>Disability</u>	<u>Discomfort</u>	<u>Satisfaction</u>	<u>Effectiveness</u>
<u>Respiratory Distress:</u>						
Patients with shortness of	<input type="radio"/> Low	<input type="radio"/> Low	<input type="radio"/> Low	<input type="radio"/> Low	<input type="radio"/> Low	<input type="radio"/> Low
breath or evidence of	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
respiratory difficulty who	<input type="radio"/> Med	<input type="radio"/> Med	<input type="radio"/> Med	<input type="radio"/> Med	<input type="radio"/> Med	<input type="radio"/> Med
continue to have spontaneous	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
breathing. May include	<input type="radio"/> High	<input type="radio"/> High	<input type="radio"/> High	<input type="radio"/> High	<input type="radio"/> High	<input type="radio"/> High
asthma, COPD, CHF.						
Excludes respiratory arrest.						

Appendix III

Measure Identification Process

Identifying measures: An example of the process for respiratory distress

Measures were identified by a process that included a literature search and structured review and discussion of the literature. The initial phase consisted of a MEDLINE search of English language articles for 1986-1996 using the Ovid (Ovid Technologies) search engine. An initial set of references was developed by combining the search for respiratory distress with a search for severity measures. Search terms for respiratory distress were developed by the investigators and consisted of: dyspnea, shortness of breath, respiration disorders, asthma, respiratory tract diseases, lung diseases, obstructive lung diseases, emphysema, reactive airways disease, croup, pulmonary edema, congestive heart failure, pneumonia, pulmonary embolism, cyanosis, anoxia, and tachypnea. Search terms for severity included: score, severity of illness index (MESH term), and predictive value of tests (MESH term). This initial set of references was then limited to articles that pertained to human subjects and published in Abridged Indexed Medicus journals.

This created a final reference set of 2,836 references. A title search of this reference list was then performed. The titles were reviewed by all of the investigators in a structured manner, identifying titles that dealt with the development or evaluation of a severity measure. Titles, which focused on a development or evaluation of the measure, were included for further review. Titles using the measures in clinical trials or for evaluations of intervention were not included. A title was included for further review if any single investigator chose it. A unanimous rejection of a title was required for the title to be eliminated from further review.

Four hundred ninety-seven (497) titles were selected for further review. Abstracts of the selected references were obtained and reviewed. The abstracts were reviewed by all the investigators. Again, an abstract required unanimous rejection by the investigators to be excluded from further evaluation. One hundred seventy-five (175) abstracts were selected for further review. For each abstract selected for

further evaluation, the full-length article was obtained. Examination of these articles resulted in 75 papers focused on the development or evaluation of a severity or outcome measure.

Articles were then sorted into groups based on the measure they were addressing (for example, dyspnea scales, or measurements of pulmonary function). A single investigator reviewed each group of articles pertaining to a specific measure. The reviews of these articles were conducted in a structured fashion. The areas considered were: time taken to complete, cost and training, scaling, and reliability. Reviews were conducted independently.

After review, the reviewers met to discuss their findings. Each reviewer presented the results of his review and made recommendations regarding the appropriateness of the measure for prehospital outcomes research. After each presentation, a discussion ensued that resulted in a consensus as to whether or not these measures should be recommended. After the measures were selected, another literature search was conducted to ensure all relevant articles regarding these measures were identified. Using the methods described above, titles, abstracts and articles were identified. These articles were then reviewed by Dr. Maio using the structured review described above. A careful review of all article references was also completed, appropriate references identified, and the complete article reviewed.

Results of the review were then discussed with the other reviewers and a consensus reached on measures to recommend. These findings were reviewed with EMSOP consultants at a project meeting. Following suggested modifications, final recommendations were made.

Appendix IV

List of Stakeholders

EMSOP Stakeholders

Agency for Health Care Research & Quality
Elinor Walker
2101 East Jefferson St., Suite 502
Rockville, MD 20852-4908
(301) 594-2049
(301) 594-2155
ewalker@ahcpr.gov

American Academy of Orthopaedic Surgeons
William W. Tipton MD, Executive Vice
President
6300 N. River Rd.
Rosemont, IL 60018
(847) 823-7186
(847) 823-8125

American Ambulance Association
Steve Haraczak, Executive Vice President
1255 23rd St, N.W.
Washington, DC 20037-1174
(202) 452-8888
(202) 452-0005
aaa911@the-aaa.org

American Association of Critical Care Nurses
Kathleen Keenan, RN, MS, CCRN, NREMT-P
101 Columbia
Aliso Viejo, CA 92656
(949) 362-2000
(949) 362-2020
info@aacn.org

American Association for Surgery of Trauma
Anthony Meyer MD
Department of Surgery
University of North Carolina
CB 7210, 167 Burnett Womack
Chapel Hill, NC 27599-7210

American College of Emergency Physicians
Colin Rorrie Jr. PhD, Executive Director
1125 Executive Circle
Irving, TX 75038-2522
(214) 550-0911
(214) 580-2816
crrorrie@acep.org

American College of Osteopathic Emergency
Physicians
Janice Wachtler, Executive Director
142 E. Ontario Street, Suite 218
Chicago, IL 60611
(312) 587-3709
(312) 587-3713

American College of Surgeons, Committee on
Trauma
Gerald Strauch MD
Director of Trauma
55 East Erie Street
Chicago, IL 60661
(312) 664-4050, x 455
(312) 440-7014
gstrauch@facs.org

American Health Information Management
Association
Mary D. Brandt
Director, Policy and Research
4610 Braeburn Dr.
Bellaire, TX 77401
(713) 668-3425
(713) 668-4973
mbrandt@ahima.mhs.compuserve.com

American Heart Association
Tom P. Aufderheide MD FACEP
Department of Emergency Medicine
Medical College of Wisconsin
8700 W. Wisconsin Ave, DH-204
Milwaukee, WI 53226
(414) 257-6060
(414) 257-8040
taufderhdi@aol.com

American Red Cross
Alice Blair
430 17th Street NW
Washington, DC 20006
(202) 639-3492
(202) 639-6116
blaira@usa.redcross.org

American Trauma Society
Harry Teter, Executive Director
8903 Presidential Parkway, Suite 512
Upper Marlboro, MD 20772
(301) 420-4189
(301) 420-0617
atstrauma@aol.com

Association for the Advancement of
Automotive Medicine
Janet Holden, Executive Director
2340 Des Plaines Avenue, Suite 106
Des Plaines, IL 60018
(847) 390-8927

Association of Air Medical Services
Dawn Mancusco, Executive Director
110 N. Royal Street, Suite 307
Alexandria, VA 22314-3234
(703) 836-8732
(703) 836-8920

Association for Health Services Research
Michael Stafford, Vice President of Operations
1130 Connecticut Avenue N.W., Suite 700
Washington, DC 20036
(202) 223-2477
(202) 835-8972
info@ahsr.org

ASTM, Committee F30 on EMS
Steve Mawn, Staff Manager
100 Barr Harbor Dr.
West Conshohocken, PA 19428-2958
(610) 832-9726
(610) 832-9666
smawn@local.astm.org

Brain Injury Association
George Zitnay
1776 Massachusetts Ave NW
Suite 100
Washington, DC 20036
(202) 296-6443
(202) 296-8850

Eastern Association of the Surgery of Trauma
Michael Rotondo MD
Brandywine Hospital
Department of Trauma
201 Reeceville Rd.
Coatsville, PA 19320

Education Development Center
Chris Miara, Manager
Children's Safety Network
55 Chapel Street
Newton, MA 02160
(617) 969-7100
(617) 244-3436

Emergency Medical Services for Children
Jane Ball, Director
Maternal and Child Health Bureau, DHHS
5600 Fishers Lane, Suite 18A-39
Rockville, MD 20857
(301) 443-4996
(301) 443-1296
jball@emscnrc.com

EMS Data Systems
Doug Brown, President
2211 E. Highland, Suite 240
Phoenix, AZ 85016
(602) 956-0126
(602) 955-5902
dbrown@emsdata.com

EMS – Research Agenda for the Future
Michael R. Sayre, M.D.
Department of Emergency Medicine
University of Cincinnati
P.O. Box 670769
231 Bethesda Avenue
Cincinnati, OH 45267-0769
(513) 558-4995
(513) 558-0378 (Fax)
Michael.Sayre@uc.edu

Lawrence Brown, EMT-P
Department of Emergency Medicine
HSC-Syracuse
750 E. Adams Street
Syracuse, NY 13210
(315) 464-4363
(315) 465-6229 (Fax)
BrownL@upstate.edu

Lynn White, MS
Department of Emergency Medicine
Akron General Medical Center
400 Wabash Avenue
Akron, OH 44307
(330) 384-6484
(330) 384-6157 (Fax)
lynn.white@worldnet.att.net

Michael Armacost, MA, NREMT-P
Colorado Department of Health
EMS Division
4300 Cherry Creek Dr., South
Denver, CO 80246-1530
(303) 692-2982
(303) 452-9396 (Fax)
mrarmaco@csn.net

J. Michael Dean, M.D., M.B.A.
Professor and Vice Chairman for Finance
Department of Pediatrics
University of Utah School of Medicine
Primary Children's Medical Center
100 North Medical Drive
Salt Lake City, Utah 84113
(801) 588-3280
(801) 588-3297 (Fax)
mike.dean@hsc.utah.edu

Scott B. Frame, M.D., FACS
Director, Division of Trauma/Critical Care
Department of Surgery
University of Cincinnati Medical Center
231 Bethesda Avenue
Cincinnati, OH 45267-0558
(513) 558-5661
(513) 558-3136 (Fax)
sbframe@prodigy.net

Baxter Larmon, PhD, MICP
Director
Center for Prehospital Care
UCLA School of Medicine
924 Westwood Blvd., #720
Los Angeles, CA 90025
(310) 794-0596
(310) 794 8796 (Fax)
blarmon@mednet.ucla.edu

Susan MacLean, RN, PhD
ENA Director of Research & Professional
Services
915 Lee Street
Des Plaines, IL 60016
(847) 460-4110
(847) 460-4005 (Fax)
smaclean@ena.org

N. Clay Mann, PhD, MS
Intermountain Injury Control Research
Center, Department of Pediatrics
University of Utah School of Medicine
410 Chipeta Way, Suite 222
Salt Lake City, Utah. 84108-1226
(801) 585-9161
(801) 581-8686 (Fax)
clay.mann@hsc.utah.edu

Gregg Margolis, MS, NREMT-P
Center for Emergency Medicine
230 McKee Place, Suite 500
Pittsburgh, PA 15213
(412) 578-3230
(412) 578-3241 (Fax)
gmargoli+@pitt.edu
Michael O'Keefe

VT Department of Health -EMS
108 Cherry Street
Burlington, VT 05402
(802) 863-7310
(802) 863-7577 (Fax)
mokeefe@vdh.state.vt.us

Daniel W. Spaite, M.D., FACEP
Emergency Medicine
1501 North Campbell Avenue
PO Box 245057
Tucson, AZ 85724-5057
(520) 626-5031
(520) 626-2480 (Fax)
dan@aemrc.arizona.edu

Jon R. Krohmer, M.D.
Kent County EMS
678 Front St., N.W., Suite 235
Grand Rapids, MI 49546
(616) 451-8438
(616) 451-8462 (Fax)
kcems1@aol.com

Emergency Nurses Association
H. Stephen Lieber, Executive Director
216 Higgins Road
Park Ridge, IL 60068
(847) 698-9400
(847) 698-9406
75250.3225@compuserve.com

Ferno
Steve Schmid
70 Weil Way
Wilmington, OH 45177
(937) 382-1451
(937) 382-1157

Health Care Financing Administration
Barbara Fleming MD PhD, Clinical Advisor
7500 Security Blvd.
Room S3-24-15, Mail Stop S2-11-07
Baltimore, MD 21244-1850
(410) 786-6863
bflaming@hcfa.gov

Health Resources and Services Administration
Jake Culp
Office of Rural Health Policy
5600 Fishers Lane, Room 9-05
Rockville, MD 20857
(301) 443-0835
(301) 443-2803
jculp@hrsa.ssw.dhhs.gov

Injury Control Research Centers (9):

1. Center for Injury Research and Control
University of Pittsburgh
230 McKee Place, Suite 400
Pittsburgh, PA 15213
<http://www.injurycontrol.com>
2. Injury Control Research Center
University of Alabama at Birmingham
403 Community Health Services Building
Birmingham, AL 35294-2041
(205) 934-7845
3. Injury Prevention Research Center
University of California at Los Angeles
School of Public Health
10833 Le Conte Avenue
Los Angeles, CA 90095-1772
(310) 206-4115
(310) 794-7989
sciprc@ucla.edu
4. Center for Injury Research and Policy
The Johns Hopkins University
School of Hygiene and Public Health
624 North Broadway
Baltimore, MD 21205
(410) 955-2636
(410) 614-2797
sogaitis@phnet.sph.jhu.edu
5. San Francisco Injury Center for
Research and Prevention
San Francisco General Hospital Building 1
San Francisco, CA 94110
(415) 821-8209
6. Harborview Injury Prevention
Research Center
Harborview Medical Center
325 9th Avenue
Box 359960
Seattle, WA 98104
(206) 521-1520
(206) 521-1562

7. Injury Prevention Research Center
University of North Carolina at Chapel
Hill
CB #7505, Chase Hall
Chapel Hill, NC 27599
(919) 966-2251
(919) 966-0466
iprc@unc.edu

8. Injury Prevention Research Center
Colorado State University
Department of Environmental Health
B107 Microbiology Building
Fort Collins, CO 80523
(970) 491-6156
(970) 491-2940

9. Injury Prevention Research Center
University of Iowa
134 AMRF, Oakdale Campus
Iowa City, IA 52242
(319) 335-4458
(319) 335-4225
john-lundell@uiowa.edu

International Association of Fire Chiefs
Jack Krakeel
4025 Fair Ridge Dr.
Fairfax, VA 22033-2868
(703) 273-0911
(703) 273-9363
krakeel@connectinc.com

International Association of Firefighters
Randall Hudgins
1750 New York Avenue, NW
Washington, DC 20006
(202) 737-8484
(202) 737-8418

Joint Commission on Accreditation of Health
Care Organizations
Deborah Nadzam PhD RN, Director,
Indicator Measurement
1 Renaissance Blvd.
Oakbrook Terrace, IL 60181
(630) 916-5240
(630) 916-5644
dnadzam@jcaho.org

Laerdal Medical Group
Al Weigel
167 Myers Corners Rd.
Wappingers Falls, NY 12590-8840
(914) 297-7770
(914) 298-4555

Medic Alert Foundation of the U.S.
Stewart Moore
2323 Colorado Avenue
Turlock, CA 95380
(209) 668-3333
(209) 669-2495

National Association of EMS Educators
Kathleen Stage-Kern
230 McKee Place, Ste. 500
Pittsburgh, PA 15213
(412) 578-3219
(412) 578-3241

National Association of EMS Physicians
Dede Gish-Panjada, Executive Director
P.O. Box 15945-281
Lenexa, KS 66285-5945
(913) 492-5858
(913) 541-0156
dpanjada@applmeapro.com

National Association of EMTs
Barbara Sanders, Executive Director
102 W. Leake St.
Clinton, MS 39056
(601) 924-7744
(601) 924-7325
namthq@aol.com

Governors Highway Safety Association
Barbara Harsha
750 First Street NE, Suite 720
Washington, DC 20002
(202) 789-0942
(202) 789-0946
naghsrb@ari.net

National Association of State EMS Directors
Elizabeth Armstrong, Executive Director
111 Park Place
Falls Church, VA 22046-4513
(703) 538-1799
(703) 241-5603
nasemsd@aol.com

National Center for Injury Prevention &
Control, CDC
Timothy B. Davis MD
4770 Buford Highway, NE (F41)
Atlanta, GA 30341-3724
(770) 488-4031
(770) 488-4338
dbd1@cdc.gov

National Committee for Quality Assurance
Richard G. Finkbiner PhD
2000 L Street NW, Suite 500
Washington, DC 20036
(202) 955-3500
(202) 955-3599
richard@ncqa.org

National Council of State EMS Training
Coordinators
Liza Burrill
Northern New Hampshire EMS,
NH Division of Health
55 Maynesboro St.
Berlin, NH 03570
(603) 752-7531
(603) 752-7533
LKBurrill@aol.com

National EMS-C Resource Alliance
James Seidel MD PhD
1124 W. Carson St., N-7
Torrance, CA 90502
(310) 222-3504
(310) 787-1763
seidelj@aol.com

National Heart, Lung and Blood Institute
Mary McDonald Hand
National Institutes of Health
31 Center Dr, MSC 2480, Room 4A16
Bethesda, MD 20892-2480
(301) 594-2726
(301) 402-1051
nadm@nhlbi.nih.gov

National Registry of EMTs
Bill Brown
6601 Busch Blvd.
P.O. Box 29233
Columbus, OH 43229
(614) 888-4484
(614) 888-8920
nremtwebocf@attmail.com

National Study Center for Trauma
Rick Bissell
701 W. Pratt St. 001
Baltimore, MD 21201
(410) 328-5085
(410) 328-3699
bissell@umbc2.umbc.edu

Orthopedic Trauma Association
Nancy Franzon, Executive Secretary
6300 N. River Rd.
Rosemont, IL 60018-4262
(847) 698-1631

Physio-Control
Robert Niskanen, Director of Research
11811 Willows Road
Redmond, WA 98073-9706
(206) 867-4437
(206) 867-4462
bniskane@physio-control.com

Society for Academic Emergency Medicine
Mary Ann Schropp, Executive Director
901 N. Washington Ave.
Lansing, MI 48906
(517) 485-5484
(517) 485-0801
faem@ix.netcom.com


Society for Trauma Nurses
Peggy Fridlund RN, MS, CEN, President
1211 Locust St.
Philadelphia, PA 19107
(215) 545-5687
(215) 545-8107
73764.123@compuserve.com

U.S. Fire Administration
Jeff T. Dyar
Office of Firefighter Safety and Health
16825 S. Seton Avenue
Emmitsburg, MD 21727
(301) 447-1333
(301) 447-1129
jdyar@fema.gov

U.S. Fire Administration
Kenneth J. Kuntz
Federal Emergency Management
Administration
NETC Building N, Room 315-C
Emmitsburg, MD 21727
(301) 447-1271
(301) 447-1102 or 447-1213
kkuntz@fema.gov

West Virginia University
Janet M. Williams MD, Research Director
Center for Rural Emergency Medicine
P.O. Box 9151
Morgantown, WV 26506-9151
(304) 293-6682
(304) 293-6702
jwillia1@wvuem1.hsc.wvu.edu

DOT HS 809 603
July 2003



U.S. Department
of Transportation
National Highway
Traffic Safety
Administration

NHTSA
People Saving People
www.nhtsa.dot.gov